

EPPO Datasheet: *Neofusicoccum laricinum*

Last updated: 2021-12-15

IDENTITY

Preferred name: *Neofusicoccum laricinum*

Authority: (Sawada) Y. Hattori & C. Nakashima

Taxonomic position: Fungi: Ascomycota: Pezizomycotina:
Dothideomycetes: Botryosphaerales: Botryosphaeriaceae

Other scientific names: *Botryosphaeria laricina* (Sawada) Shang,
Guignardia laricina (Sawada) Yamamoto & K.Ito, *Physalospora*
laricina Sawada

Common names: shoot blight of larch, twig die-back of larch

[view more common names online...](#)

EPPO Categorization: A2 list

[view more categorizations online...](#)

EU Categorization: A1 Quarantine pest (Annex II A)

EPPO Code: GUIGLA



[more photos...](#)

HOSTS


The principal hosts of *Neofusicoccum laricinum* are *Larix* spp. The most susceptible ones are *L. decidua*, *L. laricina* and *L. occidentalis*. Intermediate resistance has been observed on *L. x eurolepis* and *L. kaempferi*. Resistance has been shown on *L. gmelinii* and *L. olgensis* var. *koreana*. The only other host in nature is *Pseudotsuga menziesii*. Many other conifers can be infected by artificial inoculation. For additional information, see Sato & Shouji (1962), Sato *et al.* (1963), Ito (1963), Imazeki & Ito (1963), Oguchi (1970), Sato *et al.* (1971).

L. decidua is widely distributed in Europe at various altitudes (e.g. in the Alps and also in the Polish plains). *L. leptolepis* is also planted in the EPPO region. *Pseudotsuga menziesii* is an important forest tree.

Host list: *Larix decidua*, *Larix gmelinii* var. *japonica*, *Larix gmelinii* var. *olgensis*, *Larix gmelinii* var. *principis-ruprechtii*, *Larix gmelinii*, *Larix kaempferi*, *Larix laricina*, *Larix occidentalis*, *Larix sibirica*, *Larix x eurolepis*, *Pseudotsuga menziesii*

GEOGRAPHICAL DISTRIBUTION

N. laricinum is reported from East Asia, i.e. Eastern China, Japan, the Korean Peninsula and the Russian Far East.



EPPO Region: Russia (Far East)

Asia: China (Hebei, Heilongjiang, Jilin, Liaoning, Shandong), Japan (Hokkaido, Honshu), Korea Dem. People's Republic, Korea, Republic

BIOLOGY

The biology of *N. laricinum* has mainly been studied in Japan (Uozumi, 1961; Yokota, 1966; Sato *et al.*, 1971).

The asexual morph appears in abundance on the underside of needles and on young sprouts between July and November, and spores are dispersed by insects or rain. During this time, the pycnidiospores are released and give rise to secondary infections in late summer. Discharge of conidia occurs between 10 and 35°C (25°C optimum) and was observed to occur at 98% RH. A few spores in their pycnidia can overwinter until the following April.

The sexual morph appears on branches after October. The black pseudothecia, which occur in groups or singly, take 2 years to develop. Ascospores released between May and October (peak July-August) are the source of primary infections. Optimum temperature for infection is 20°C with free water. Ascospores can infect host plants throughout the season, but do so principally at the beginning of August; wounds do not appear necessary for penetration. Disease symptoms appear about 2 weeks after infection. Some spores may overwinter in the pseudothecia. Cool winters and short summers do not favour the disease.

DETECTION AND IDENTIFICATION

Symptoms

The disease is conspicuous as discoloration, wilting and death of the succulent current season's growth. Old twigs remain unaffected. Early attack, visible between June and September, causes hanging-down of the top of shoots, accompanied by a yellowing and browning of needles which may fall. The needles at the tops of shoots turn brown and often remain on the tree during winter. Dark, sunken lesions, abundant in sporulating bodies, and exuding resin appear on the stems of affected seedlings and on shoots, and usually girdle these parts. The resin hardens into whitish drops. Late infections, occurring in September to early October, do not show the characteristic hanging-down, owing to the lignified nature of the twigs. On needles, symptoms appear as brown spots with chlorotic haloes, which subsequently coalesce. Repeated infections result in stunted, bushy trees with many dead shoots.

For additional information, see Imazeki & Ito (1963), Ito (1963), Sato *et al.* (1971).

Morphology

Sexual morph: Diseased twigs defoliated from the middle to the tip, with exudate resin. Fruit bodies lined, erumpent. Ascomata epidermal, blackish, globose, 368 μm diam; ostiole erumpent, 60 μm diam; paraphyses developed, intricate, 3 μm . Ascus clavate, rounded at the apex, stipitate at the base, hyaline, 114–135 \times 22–26 μm . Ascospores ellipsoid, smooth, hyaline, 24–27 \times 13 μm (Sawada, 1950).

Asexual morph: Conidiomata pycnidial, epidermal, merged, solitary, globose, dark brown, subglobose, unilocular, with a central ostiole, 204–246 \times 207–212 μm ; pycnidial wall composed of depressed or irregular cells in three to four layers, brown to dark brown. Conidiophores reduced to conidiogenous cells; conidiogenous cells discrete, hyaline, cylindrical to ampulliform, determinate, with periclinal thickening, or proliferating percurrently, 9–23 \times 2.4–5 μm . Paraphyses not seen. Conidia holoblastic sporulation for first conidia, phialidic sporulation for following conidia, hyaline, smooth, aseptate, slightly colored and septate with age, ellipsoid to fusiform, granulate, subtruncate to bluntly rounded at the base, rounded to subacute at the apex, with a short frill at both ends, 23–38 \times 7–12 μm , 29.85 \times 8.50 μm on average, L/W = 3.57 (Hattori *et al.* 2021).

Detection and inspection methods

The fungal fruiting bodies may be observed directly or isolated and cultured on a medium containing 3 g yeast extract, 10 g soluble starch, 0.25 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g agar in 1 L distilled water, maintained at 20°C (Hara & Ito, 1963; Ito, 1963). To identify species in the genus *Neofusicoccum*, it is now essential to use molecular phylogenetic analysis using the regions of rDNA ITS, rpb2, tef1-?, and tub2. The deposit numbers for each ex-epitype (FFPRI 411215 = MUCC 2662) sequence are LC589129 for ITS, LC589140 for tef1, LC589151 for tub2, LC589164 for rpb2 (Hattori *et al.*, 2021).

PATHWAYS FOR MOVEMENT

Under natural conditions, *N. laricinum* spreads by dispersal of ascospores and conidia. In international trade, spread is possible on diseased host trees, including artificially dwarfed plants. Cut branches may also be a possible pathway (EFSA, 2018). Pollen or seed is unlikely to harbour the pathogen.

PEST SIGNIFICANCE

Economic impact

N. laricinum caused the most serious disease of *Larix* forests and nurseries in Japan. It had long been known locally, but started to cause large-scale damage in *Larix* plantations after 1959 - at which time, areas planted with *Larix* were increasing rapidly. In 1963, more than 80 000 ha of plantations were diseased, with 100% of the trees affected. Although young diseased trees do not usually die, their subsequent growth is retarded or stopped. *N. laricinum* causes severe damage in areas with strong winds, but after changing the tree species planted in these areas and using appropriate management measures, this disease is now under control.

Control

In Japan, chemical control is applied against this disease (Oguchi, 1980; Okada, 2000). In addition, intensive testing of disease-resistant clones and observation of their growth in forest plantations has been carried out (Oguchi, 1963; Sato, 1970; Kobayashi, 1980).

In nursery fields, fungicides can be applied every 2 weeks during the infection period (June to September). In highly infested nurseries, dipping of *Larix* seedlings into fungicide solution in spring is also used. In forest plantations, it is important to avoid bringing in diseased seedlings. When the disease is not widespread, early detection is important to eradicate the disease. If damage is widespread, it is necessary to reduce the density of the pathogen by prioritizing

the felling of diseased trees during thinning. Removal and burning of the infected trees and reforestation by other species are also carried out in heavily diseased stands.

Phytosanitary risk

In the EPPO region, *N. laricinum* could be potentially dangerous to *Larix* and *P. menziesii*, wherever present. Considering the distribution of *N. laricinum* in East Asia, climatic conditions prevailing in Europe are assumed not to be a limiting factor (EFSA, 2018).

PHYTOSANITARY MEASURES

Considering the risk that this fungus could present to the EPPO region it is recommended that all countries should prohibit importation of plants for planting and cut branches of *Larix* and *P. menziesii* from countries where *N. laricinum* this fungus occurs.

REFERENCES

EFSA PLH Panel (EFSA Panel on Plant Health) Jeger M, Bragard C, Caffier D, Candresse T, Chatzivassiliou E, Dehnen-Schmutz K, Gilioli G, Gregoire J-C, Jaques Miret JA, MacLeod A, Navajas Navarro M, Niere B, Parnell S, Potting R, Rafoss T, Rossi V, Urek G, Van Bruggen A, Van der Werf W, West J, Winter S, Boberg J, Gonthier P, Pautasso M (2018) Scientific Opinion on the pest categorisation of *Guignardia laricina*. *EFSA Journal* **16**(6), 5303, 24 pp. <https://doi.org/10.2903/j.efsa.2018.5303>

Hara K & Ito K (1963) [Agar-media for sporulation of *Guignardia laricina* (Sawada) W. Yamamoto et K. ITO, the shoot blight fungus of larch (Preliminary report)]. *Journal of the Japanese Forest Society* **45**, 238-241 (In Japanese)

Hattori Y, Ando Y & Nakashima C (2021) Taxonomical re-examination of the genus *Neofusicoccum* in Japan. *Mycoscience* **62**, 250-259.

Imazeki R & Ito K (1963) Dangerous forest diseases in Japan. Shoot blight of larch. In: *Internationally dangerous forest tree diseases. Miscellaneous Publications, Forest Service, US Department of Agriculture* No. 939, 48-49.

Ito K (1963) Shoot blight of larches. A destructive disease in larch plantations of Japan. *Bulletin of the Government Forest Experiment Station, Tokyo* **159**, 89-103.

Kobayashi T (1980) Important forest diseases and their control measures. *Plant Protection in Japan, Agriculture Asia*, Special Issue No. 11, 298 pp.

Okada M (2000) [The shoot blight disease of Larch], [?], No. 103 (in Japanese).

Oguchi T (1963) [Differential resistance of Larch clones to the shoot blight of Larch], [?], NO. 2 (in Japanese).

Oguchi T (1970) Growth of three species of larch and their infecting period to the shoot blight disease. *Bulletin, Hokkaido Forest Experiment Station* **8**, 35-40.

Oguchi T (1980) [Forest Diseases Series 1: Four Major Diseases of Japanese Larch]. [?], 44, 46 (in Japanese).

Sato K & Shouji T (1962) [Ditto V (Preliminary report). Pathogenicity of *Guignardia laricina*, the causal fungus of the disease]. *Transactions of the 73rd Annual Meetings of the Japanese Forestry Society*, 217-219 (In Japanese).

Sato K, Yokozawa Y, Shoji T (1963) Studies on the shoot blight disease of larch I. *Bulletin of the Government Forest Experimental Station, Tokyo* **156**, 85-137.

Sato K (1970) [Disease resistance of silvicultural tree species and their cultivars in the

Tohoku region]. ??????, No. 26 (in Japanese).

Sato K, Yoshinori Y, Shoji T, Kojima C (1971) Studies on the shoot blight of larch II. *Bulletin of the Government Forest Experiment Station, Meguro* **236**, 27-91.

Sawada (1950) Fungi inhabiting conifers in the Tohoku district. II. Fungi on various conifers except 'Sugi'. *Bulletin of the Government Forest Experimental Station, Meguro* **46**, 144–148.

Uozumi T (1961) Studies on the shoot blight disease of larch. With special reference to life history of the causal fungus, *Physalospora laricina* Sawada. *Bulletin of the Government Forest Experiment Station, Meguro* **132**, 47-54.

Yokota S (1966) Ecological studies on *Guignardia laricina*, the causal fungus of the shoot blight of larch trees and climatic factors influencing the outbreak of the disease. *Bulletin of the Government Forest Experiment Station, Meguro* **184**, 79 pp.

ACKNOWLEDGEMENTS

This datasheet was extensively revised in 2021 by Yukako Hattori, Forestry and Forest Products Research Institute (FFPRI), Japan Society for the Promotion of Science (JSPS). Her valuable contribution is gratefully acknowledged.

How to cite this datasheet?

EPPO (2024) *Neofusicoccum laricinum*. EPPO datasheets on pests recommended for regulation. Available online. <https://gd.eppo.int>

Datasheet history

This datasheet was first published in the EPPO Bulletin in 1978, revised in the two editions of 'Quarantine Pests for Europe' in 1992 and 1997, as well as in 2021. It is now maintained in an electronic format in the EPPO Global Database. The sections on 'Identity', 'Hosts', and 'Geographical distribution' are automatically updated from the database. For other sections, the date of last revision is indicated on the right.

CABI/EPPO (1992/1997) *Quarantine Pests for Europe (1st and 2nd edition)*. CABI, Wallingford (GB).

EPPO (1978) Data sheets on quarantine organisms No. 12, *Guignardia laricina*. *EPPO Bulletin* **8**(2), 2 pp. <https://doi.org/10.1111/j.1365-2338.1978.tb02759.x>



Co-funded by the
European Union