

EPPO Datasheet: *Orthospovirus tomatomaculae*

Last updated: 2021-08-30

IDENTITY

Preferred name: *Orthospovirus tomatomaculae*

Taxonomic position: Viruses and viroids: Riboviria: Orthornavirae: Negarnaviricota: Polyploviricotina: Bunyaviricetes: Elliovirales: Tospoviridae

Other scientific names: TSWV, Tomato spotted wilt orthospovirus, Tomato spotted wilt tospovirus, Tomato spotted wilt virus

Common names: bronze leaf of tomato, kromnek virus, spotted wilt of tomato, yellow spot of pineapple

[view more common names online...](#)

EPPO Categorization: A2 list

[view more categorizations online...](#)

EU Categorization: RNQP (Annex IV)

EPPO Code: TSWV00



[more photos...](#)

Notes on taxonomy and nomenclature

Tomato spotted wilt disease was first described in Australia in 1915 and was later identified as a viral disease caused by tomato spotted wilt virus (TSWV) (Samuel *et al.*, 1930). In 1990, TSWV was assigned to the genus *Tospovirus*, which was later renamed *Orthospovirus* (family *Tospoviridae*, order *Bunyavirales*; ICTV online: <https://talk.ictvonline.org/taxonomy/>; Abudurexiti *et al.*, 2019). Species demarcation within the genus is based on nucleoprotein (N) sequence (new species are defined as having less than 90% amino acid sequence similarity to all other described species within the genus); in addition, species are often biologically distinguished by their host range and vector specificity (Plyusnin *et al.*, 2011; Kormelink *et al.*, 2021).

HOSTS

TSWV has one of the largest known host ranges of any plant virus, with more than 1000 plant species recorded as hosts. The host range includes dicotyledons and monocotyledons, crop plants, ornamentals and weeds. Most of the plant species susceptible to TSWV belong to the *Asteraceae* and *Solanaceae* families (Parrella *et al.*, 2003).

In the EPPO region, the main crop hosts are artichokes (*Cynara scolymus*), eggplants (*Solanum melongena*), peppers (*Capsicum annuum*), chicory (*Cichorium* spp.), cucurbits (*Cucurbitaceae*), faba beans (*Vicia faba*), lettuce (*Lactuca sativa*), potatoes (*Solanum tuberosum*), tobacco (*Nicotiana tabacum*), tomatoes (*Solanum lycopersicum*); the main ornamental hosts are *Anemone*, *Aster*, *Begonia* hybrids, *Calceolaria*, *Callistephus*, chrysanthemum (*Dendranthema x grandiflorum*), *Cyclamen*, *Dahlia*, *Gerbera*, *Pelargonium*, *Pericallis cruenta*, *Ranunculus*, *Sinningia* spp., *Symphytichum*, *Tagetes patula* and *Zinnia*. Wild species such as *Senecio vulgaris*, *Solanum nigrum*, *Sonchus* spp. and *Stellaria media* can be important reservoirs for TSWV.

Host list: *Acalypha australis*, *Acanthospermum hispidum*, *Acanthus mollis*, *Achimenes* sp., *Aconitum carmichaelii*, *Adenium obesum*, *Aegopodium podagraria*, *Aeschynanthus pulcher*, *Aeschynanthus speciosus*, *Agapanthus praecox* subsp. *orientalis*, *Agapanthus praecox*, *Agastache foeniculum*, *Ageratum houstonianum*, *Aglaonema commutatum*, *Agrostemma githago*, *Ajania pacifica*, *Alcea rosea*, *Alkekengi officinarum*, *Alliaria petiolata*, *Allium ascalonicum*, *Allium cepa*, *Allium porrum*, *Allium sativum*, *Alstroemeria aurea*, *Alstroemeria* hybrids, *Alternanthera sessilis*, *Althaea officinalis*, *Amaranthus albus*, *Amaranthus blitum*, *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus graecizans*, *Amaranthus hybridus*, *Amaranthus palmeri*, *Amaranthus powellii*, *Amaranthus retroflexus*, *Amaranthus spinosus*, *Amaranthus thunbergii*, *Amaranthus viridis*, *Amaryllis belladonna*, *Ambrosia artemisiifolia*, *Ambrosia psilostachya*, *Ambrosia trifida*, *Ammi majus*, *Ananas comosus*, *Anchusa azurea*, *Anchusa capensis*, *Anchusa officinalis*, *Anchusa undulata*, *Anemone coronaria*, *Antennaria neglecta*, *Anthemis arvensis*, *Anthurium andraeanum*, *Anthurium hookeri*, *Anthurium scherzerianum*, *Antirrhinum majus*, *Aphelandra squarrosa*, *Apium graveolens* var. *dulce*, *Apium graveolens*, *Aquilegia vulgaris*, *Arabis* sp., *Arachis batizocoi*, *Arachis duranensis*, *Arachis hypogaea*


, *Arachis monticola*, *Arachis pusilla*, *Arachis stenosperma*, *Arachis villosulicarpa*, *Arctium lappa*, *Arctium minus*, *Arctotheca calendula*, *Arctotis x hybrida*, *Ardisia sp.*, *Argyranthemum frutescens*, *Aristolochia clematitidis*, *Aristolochia elegans*, *Artemisia absinthium*, *Artemisia dracunculus*, *Artemisia princeps*, *Arum maculatum*, *Arum palaestinum*, *Asarum canadense*, *Asclepias curassavica*, *Asparagus setaceus*, *Asplenium nidus*, *Aster amellus*, *Aubrieta deltoidea*, *Avena fatua*, *Ballota nigra*, *Barbarea vulgaris*, *Begonia gracilis*, *Begonia semperflorens hybrids*, *Begonia tuberhybrida hybrids*, *Begonia x hiemalis*, *Bellis caerulea*, *Berberis sp.*, *Beta vulgaris*, *Betonica officinalis*, *Bidens discoidea*, *Bidens pilosa*, *Bidens subalternans*, *Bidens vulgata*, *Blitum capitatum*, *Blitum virgatum*, *Bothriocline longipes*, *Bouvardia sp.*, *Brachyscome iberidifolia*, *Brassica juncea*, *Brassica napus*, *Brassica oleracea var. botrytis*, *Brassica oleracea*, *Brassica perviridis*, *Brassica rapa subsp. chinensis*, *Brassica rapa subsp. pekinensis*, *Brassica rapa subsp. sylvestris*, *Browallia americana*, *Browallia speciosa*, *Brugmansia arborea*, *Brugmansia aurea*, *Brugmansia suaveolens*, *Buddleia davidii*, *Cajanus cajan*, *Calceolaria crenatiflora*, *Calceolaria herbeohybrida hybrids*, *Calectasia cyanea*, *Calendula arvensis*, *Calendula officinalis*, *Calla palustris*, *Callistephus chinensis*, *Calotropis gigantea*, *Calystegia sepium*, *Campanula americana*, *Campanula glomerata*, *Campanula grandis subsp. grandis*, *Campanula isophylla*, *Campanula medium*, *Campanula persicifolia*, *Campanula pyramidalis*, *Campanula rapunculoides*, *Canavalia ensiformis*, *Canavalia gladiata*, *Canna sp.*, *Capsella bursa-pastoris*, *Capsicum annuum*, *Capsicum chinense*, *Capsicum frutescens*, *Cardamine flexuosa*, *Cardamine hirsuta*, *Cardamine oligosperma*, *Cardamine parviflora*, *Carduus acanthoides*, *Carduus nutans*, *Carica papaya*, *Carthamus tinctorius*, *Carum carvi*, *Catharanthus roseus*, *Celosia argentea*, *Centaurea cineraria*, *Centaurea cyanus*, *Centella asiatica*, *Centranthus ruber*, *Cerastium glomeratum*, *Cerastium holosteoides*, *Cestrum aurantiacum*, *Cestrum elegans*, *Cestrum nocturnum*, *Cestrum parqui*, *Chaerophyllum temulum*, *Chamaedorea costaricana*, *Chamaedorea elegans*, *Chelone sp.*, *Chenopodium murale*, *Chenopodium simplex*, *Chenopodium album*, *Chenopodium ficifolium*, *Chenopodium giganteum*, *Chenopodium vulvaria*, *Chondrilla juncea*, *Chrysanthemum indicum*, *Chrysanthemum x morifolium*, *Cicer arietinum*, *Cichorium endivia var. latifolia*, *Cichorium endivia*, *Cichorium intybus*, *Cineraria sp.*, *Cirsium arvense*, *Cirsium japonicum*, *Cirsium palustre*, *Cirsium setidens*, *Cirsium vulgare*, *Citrullus lanatus*, *Clarkia amoena subsp. lindleyi*, *Claytonia perfoliata*, *Clematis flammula*, *Clematis vitalba*, *Clerodendrum thomsoniae*, *Clivia sp.*, *Codonanthe crassifolia*, *Codonopsis pilosula*, *Coffea arabica*, *Coleus australis*, *Coleus scutellarioides*, *Columnea hirta*, *Commelina communis*, *Commelina cyanea*, *Conium maculatum*, *Convolvulus althaeoides*, *Convolvulus arvensis*, *Convolvulus cantabrica*, *Coprosma repens*, *Cordyline fruticosa*, *Coreopsis basalis*, *Coreopsis lanceolata*, *Coreopsis tinctoria*, *Coriandrum sativum*, *Cornus sanguinea*, *Cosmos bipinnatus*, *Cota tinctoria*, *Crepis capillaris*, *Crepis divaricata*, *Crepis foetida subsp. rhoeadifolia*, *Crepis occidentalis subsp. pumila*, *Crepis pulchra*, *Crinum jagus*, *Crinum moorei*, *Crotalaria incana*, *Crotalaria juncea*, *Crotalaria pallida var. obovata*, *Crotalaria spectabilis*, *Croton glandulosus*, *Cucumis melo*, *Cucumis sativus*, *Cucurbita maxima*, *Cucurbita moschata*, *Cucurbita pepo*, *Cyanus montanus*, *Cyclamen persicum*, *Cymbidium sp.*, *Cynanchum rostellatum*, *Cynara cardunculus*, *Cynara scolymus*, *Cynodon dactylon*, *Cynoglossum sp.*, *Cyperus esculentus*, *Cyperus rotundus*, *Cyrtanthus sp.*, *Cytisus scoparius*, *Dahlia hybrids*, *Dahlia pinnata*, *Datura ferox*, *Datura innoxia*, *Datura metel*, *Datura stramonium var. tatula*, *Datura stramonium*, *Datura wrightii*, *Delphinium hybrids*, *Dendrobium sp.*, *Desmodium tortuosum*, *Desmodium uncinatum*, *Dianthus chinensis*, *Diascia barberae*, *Dieffenbachia sp.*, *Digitalis sp.*, *Digitaria sanguinalis*, *Dimorphotheca ecklonis*, *Dimorphotheca fruticosa*, *Dimorphotheca pluvialis*, *Dimorphotheca sinuata*, *Diploaxis erucoides*, *Diploaxis muralis*, *Dorotheanthus tricolor*, *Dracaena draco*, *Dracaena fragrans*, *Dracaena marginata*, *Duboisia leichhardtii*, *Duboisia myoporoides*, *Dysphania ambrosioides*, *Echinacea purpurea*, *Echinochloa colonum*, *Echinocystis lobata*, *Eclipta prostrata*, *Eleusine indica*, *Emilia coccinea*, *Emilia sonchifolia*, *Epilobium hirsutum*, *Epipremnum pinnatum*, *Erigeron bonariensis*, *Erigeron canadensis*, *Erigeron strigosus*, *Erodium ciconium*, *Erodium moschatum*, *Erysimum cheiranthoides*, *Erysimum x cheiri*, *Eupatorium capillifolium*, *Eupatorium maculatum*, *Euphorbia heterophylla*, *Euphorbia peplus*, *Euphorbia pulcherrima*, *Euphorbia tithymaloides*, *Euryops sp.*, *Eustoma russellianum*, *Exacum affine*, *Fagopyrum esculentum*, *Fallopia convolvulus*, *Farfugium japonicum*, *Fatsyhedera lizei*, *Fatsia japonica*, *Felicia amelloides*, *Felicia bergeriana*, *Felicia fruticosa*, *Ficus benjamina*, *Ficus cyathistipula*, *Ficus elastica*, *Ficus lyrata*, *Flaveria bidentis*, *Forsythia koreana*, *Fragaria vesca*, *Freesia hybrids*, *Fuchsia hybrids*, *Fumaria muralis*, *Fumaria officinalis*, *Gaillardia aristata*, *Gaillardia x grandiflora*, *Galinsoga parviflora*, *Galinsoga quadriradiata*, *Galium aparine*, *Galium lucidum*, *Galium parisiense*, *Galium spurium*, *Galium tricornutum*, *Galium verum*, *Gamochaeta falcata*, *Gamochaeta pennsylvanica*, *Gamochaeta purpurea*, *Gardenia jasminoides*, *Gazania rigens*, *Geranium carolinianum*, *Geranium robertianum*, *Geranium rotundifolium*, *Gerbera jamesonii*, *Gerbera x hybrida*, *Gesneria sp.*, *Gladiolus grandiflorus*, *Glaucium flavum*, *Glebionis coronaria*, *Gleditsia triacanthos*, *Gloriosa superba*, *Gloxinia sp.*, *Glycine max*, *Glycine soja*, *Gnaphalium uliginosum*, *Gomphrena globosa*, *Goniolimon tataricum*, *Gossypium hirsutum*, *Guzmania sp.*, *Gymnema sylvestre*, *Gynura aurantiaca*, *Gypsophila elegans*, *Gypsophila paniculata*, *Helianthus annuus*, *Helichrysum sp.*, *Heliotropium europaeum*, *Helminthotheca echioides*, *Hemistepta lyrata*, *Heptapleurum actinophyllum*, *Heptapleurum arboricola*, *Hesperis matronalis*, *Hibiscus trionum*, *Hippeastrum aulicum*, *Hippeastrum hybrids*, *Hippeastrum reginae*, *Hippeastrum striatum*, *Hirschfeldia incana*, *Holcus lanatus*

, *Hoya australis*, *Hoya bella*, *Hoya carnosae*, *Hoya linearis*, *Humulus scandens*, *Hydrangea macrophylla*, *Hymenocallis occidentalis*, *Hyoscyamus niger*, *Hypericum triquetrifolium*, *Iberis semperflorens*, *Impatiens New Guinea hybrids*, *Impatiens balsamina*, *Impatiens capensis*, *Impatiens hybrids*, *Impatiens walleriana*, *Inula helenium*, *Ipomoea batatas*, *Ipomoea indica*, *Ipomoea lacunosa*, *Ipomoea purpurea*, *Iris domestica*, *Iris ensata*, *Iris x hollandica*, *Jacobaea maritima*, *Jacobaea vulgaris*, *Jacquemontia tamnifolia*, *Jasminum odoratissimum*, *Justicia brandegeana*, *Kalanchoe blossfeldiana*, *Kalanchoe daigremontiana*, *Kalanchoe synsepala*, *Kalanchoe thyrsiflora*, *Lablab purpureus*, *Lactuca canadensis*, *Lactuca floridana*, *Lactuca indica*, *Lactuca saligna*, *Lactuca sativa var. capitata*, *Lactuca sativa var. crispata*, *Lactuca sativa var. longifolia*, *Lactuca sativa*, *Lactuca serriola*, *Lagascea mollis*, *Lagenaria siceraria*, *Lamium amplexicaule*, *Lamium purpureum*, *Lantana camara*, *Lathyrus odoratus*, *Lathyrus sativus*, *Lavandula angustifolia subsp. pyrenaica*, *Layia elegans*, *Leonotis nepetifolia*, *Lepidium didymum*, *Lepidium draba*, *Lepidium sativum*, *Lepidium virginicum*, *Leptochiton quitoensis*, *Leucanthemum maximum*, *Leucanthemum vulgare*, *Leucanthemum x superbum*, *Ligustrum vulgare*, *Lilium lancifolium*, *Lilium longiflorum*, *Limonium perezii*, *Limonium platyphyllum*, *Limonium sinuatum*, *Limonium vulgare*, *Lisianthus sp.*, *Lobelia dortmanna*, *Lobelia erinus*, *Lobelia valida*, *Lobularia maritima*, *Lolium perenne*, *Luffa aegyptiaca*, *Lupinus albus*, *Lupinus angustifolius*, *Lupinus leucophyllus*, *Lupinus polyphyllus*, *Lupinus subcarnosus*, *Lychnis chalcidonica*, *Lycianthes rantonnetii*, *Lycium chinense*, *Lycium ferocissimum*, *Lycopus europaeus*, *Lysimachia arvensis*, *Lysimachia congestiflora*, *Lysimachia loeflingii*, *Lysimachia nummularia*, *Macrotyloma uniflorum*, *Maianthemum canadense*, *Malcolmia maritima*, *Malva neglecta*, *Malva nicaeensis*, *Malva parviflora*, *Malva pusilla*, *Malva sylvestris*, *Malva verticillata*, *Maranta leuconeura*, *Marrubium vulgare*, *Martynia annua*, *Matricaria chamomilla*, *Matricaria discoidea*, *Matthiola incana*, *Mazus pumilus*, *Medicago lupulina*, *Medicago polymorpha*, *Medinilla magnifica*, *Melampodium sp.*, *Melilotus indicus*, *Melilotus officinalis*, *Melissa officinalis*, *Mentha arvensis*, *Mentha microphylla*, *Mentha spicata*, *Mentha suaveolens*, *Mentha x piperita*, *Mercurialis annua*, *Mimulus sp.*, *Mirabilis jalapa*, *Mollugo verticillata*, *Moluccella laevis*, *Monarda didyma*, *Monarda fistulosa*, *Montia sp.*, *Morus alba*, *Myosotis alpestris*, *Myosoton aquaticum*, *Nasturtium officinale*, *Nepeta cataria*, *Nepeta nuda*, *Nerine bowdenii*, *Nerium oleander*, *Nicandra physalodes*, *Nicotiana acuminata*, *Nicotiana alata*, *Nicotiana benthamiana*, *Nicotiana bonariensis*, *Nicotiana debneyi*, *Nicotiana excelsior*, *Nicotiana exigua*, *Nicotiana glauca*, *Nicotiana goodspeedii*, *Nicotiana langsdorffii*, *Nicotiana longiflora*, *Nicotiana maritima*, *Nicotiana megalosiphon*, *Nicotiana nudicaulis*, *Nicotiana paniculata*, *Nicotiana pauciflora*, *Nicotiana plumbaginifolia*, *Nicotiana quadrivalvis*, *Nicotiana solanifolia*, *Nicotiana suaveolens*, *Nicotiana sylvestris*, *Nicotiana tabacum*, *Nicotiana tomentosiformis*, *Nicotiana undulata*, *Nicotiana velutina*, *Nicotiana wigandioides*, *Nicotiana x edwardsonii*, *Nicotiana x sanderi*, *Nolana sp.*, *Notobasis syriaca*, *Nuttallanthus canadensis*, *Ocimum basilicum*, *Oenanthe javanica*, *Oenothera biennis*, *Oenothera laciniata*, *Oenothera sp.*, *Oncidium sp.*, *Onopordum acanthium*, *Onopordum illyricum*, *Opuntia sp.*, *Orlaya sp.*, *Ornithogalum thyrsoides*, *Orobanche sp.*, *Osteospermum sp.*, *Oxalis acetosella*, *Oxalis corniculata*, *Oxalis latifolia*, *Oxalis stricta*, *Oxalis tuberosa*, *Oxybasis glauca*, *Oxybasis urbica*, *Oxypetalum sp.*, *Pachypodium lamerei*, *Paederia foetida*, *Paeonia lactiflora*, *Papaver dubium*, *Papaver nudicaule*, *Papaver orientale*, *Papaver rhoeas*, *Papaver somniferum*, *Parietaria officinalis*, *Pelargonium crispum*, *Pelargonium grandiflorum*, *Pelargonium peltatum*, *Pelargonium x hortorum*, *Pelargonium zonale*, *Penstemon hartwegii*, *Penstemon hirsutus*, *Peperomia fraseri*, *Peperomia obtusifolia*, *Peperomia rotundifolia*, *Pericallis cruenta*, *Pericallis x hybrida*, *Persicaria hydropiper*, *Persicaria lapathifolia*, *Persicaria longiseta*, *Persicaria maculosa*, *Persicaria pennsylvanica*, *Petasites japonicus*, *Petroselinum crispum*, *Petunia hybrids*, *Petunia integrifolia*, *Phalaenopsis sp.*, *Phaseolus lunatus*, *Phaseolus vulgaris*, *Philodendron sagittifolium*, *Phlox drummondii*, *Physalis acutifolia*, *Physalis alkekengi var. franchetii*, *Physalis angulata*, *Physalis heterophylla*, *Physalis ixocarpa*, *Physalis minima*, *Physalis peruviana*, *Physalis pruinosa*, *Physalis pubescens*, *Physostegia virginiana*, *Phytolacca americana*, *Pilea pumila*, *Pinus sp.*, *Pisum sativum subsp. arvense*, *Pisum sativum*, *Pittosporum tobira*, *Plantago lanceolata*, *Plantago major*, *Plantago rugelii*, *Platycerium superbum*, *Platycodon grandiflorus*, *Plumeria rubra f. acutifolia*, *Poa annua*, *Polygonum aviculare*, *Portulaca grandiflora*, *Portulaca oleracea*, *Portulaca pilosa*, *Potentilla reptans*, *Primula malacoides*, *Primula obconica*, *Primula sinensis*, *Primula vulgaris*, *Prunella vulgaris*, *Pyrrhopappus carolinianus*, *Ranunculus abortivus*, *Ranunculus acris*, *Ranunculus arvensis*, *Ranunculus asiaticus*, *Ranunculus bulbosus*, *Ranunculus hybridus*, *Ranunculus muricatus*, *Ranunculus sardous*, *Raphanus raphanistrum*, *Raphanus sativus*, *Rhamnus sp.*, *Rhaponticum carthamoides*, *Rhodanthe chlorocephala subsp. rosea*, *Rhodanthe manglesii*, *Rhododendron sp.*, *Richardia scabra*, *Ricinus communis*, *Robinia pseudoacacia*, *Rohdea sp.*, *Rorippa indica*, *Rorippa palustris*, *Rosa sp.*, *Rubus idaeus*, *Rubus ulmifolius*, *Rudbeckia amplexicaulis*, *Rudbeckia hirta*, *Rudbeckia nitida*, *Rudbeckia occidentalis*, *Rumex conglomeratus*, *Rumex crispus*, *Ruscus hypoglossum*, *Salpiglossis sinuata*, *Salsola kali subsp. ruthenica*, *Salvia officinalis*, *Salvia pratensis*, *Salvia sclarea*, *Salvia splendens*, *Salvia verbenaca*, *Sambucus nigra*, *Sanguisorba minor*, *Saponaria officinalis*, *Saxifraga stolonifera*, *Scabiosa atropurpurea*, *Schizanthus pinnatus*, *Schlumbergera truncata*, *Schlumbergera x buckleyi*, *Scutellaria sp.*, *Sechium edule*, *Sedum sarmentosum*, *Seemannia sylvatica*, *Senecio doria*, *Senecio vulgaris*, *Senna obtusifolia*, *Senna occidentalis*, *Senna tora*, *Sesamum indicum*, *Setaria viridis*, *Sida rhombifolia*, *Sida spinosa*, *Sigesbeckia orientalis*, *Silene coronaria*, *Silene firma*, *Silene latifolia subsp. alba*, *Silene vulgaris*

, *Sinapis arvensis*, *Sinningia speciosa*, *Sisymbrium irio*, *Smilax sieboldii*, *Solanum aculeatissimum*, *Solanum americanum*, *Solanum aviculare*, *Solanum betaceum*, *Solanum capsicoides*, *Solanum carolinense*, *Solanum chenopodioides*, *Solanum chilense*, *Solanum dulcamara*, *Solanum grandiflorum*, *Solanum habrochaites*, *Solanum hieronymi*, *Solanum laciniatum*, *Solanum lycocarpum*, *Solanum lycopersicum*, *Solanum mammosum*, *Solanum marginatum*, *Solanum melongena*, *Solanum muricatum*, *Solanum neorickii*, *Solanum nigrum*, *Solanum palinacanthum*, *Solanum pennellii*, *Solanum pimpinellifolium*, *Solanum pseudocapsicum*, *Solanum quitoense*, *Solanum robustum*, *Solanum seaforthianum*, *Solanum triflorum*, *Solanum trifolium*, *Solanum tuberosum*, *Solanum tweedeanum*, *Solanum viarum*, *Solanum villosum* subsp. *miniatum*, *Solanum violaceum*, *Solidago altissima*, *Solidago canadensis*, *Sonchus asper*, *Sonchus oleraceus*, *Sonchus tenerrimus*, *Sorbus aucuparia*, *Spathiphyllum floribundum*, *Spergula arvensis*, *Spinacia oleracea*, *Stachys arvensis*, *Stapelia* sp., *Stellaria media*, *Stellaria uliginosa*, *Stephanotis floribunda*, *Stevia rebaudiana*, *Streptocarpus ionanthus*, *Streptosolen jamesonii*, *Symphoricarpos albus*, *Symphyotrichum cordifolium*, *Symphyotrichum ericoides*, *Symphyotrichum lateriflorum*, *Symphyotrichum novae-angliae*, *Symphytum tuberosum*, *Syngonium podophyllum*, *Tagetes erecta*, *Tagetes minuta*, *Tagetes patula*, *Taraxacum officinale*, *Tephrosia purpurea*, *Tetragonia tetragonioides*, *Thymus vulgaris*, *Tithonia rotundifolia*, *Tolmiea menziesii*, *Torilis arvensis*, *Trachelium caeruleum*, *Trachymene coerulea*, *Tradescantia albiflora*, *Tragopogon dubius*, *Tragopogon porrifolius*, *Tragopogon pratensis*, *Tragopogon x mirus*, *Tragopogon x miscellus*, *Trevesia palmata*, *Tribulus terrestris*, *Trichosanthes kirilowii*, *Trifolium incarnatum*, *Trifolium repens*, *Trifolium* sp., *Trifolium subterraneum*, *Trifolium tembense*, *Triodanis perfoliata*, *Tripleurospermum inodorum*, *Tropaeolum majus*, *Tulbaghia violacea*, *Tussilago farfara*, *Urceolina x grandiflora*, *Urtica dioica*, *Valeriana fauriei*, *Valeriana officinalis*, *Valerianella locusta*, *Verbascum blattaria*, *Verbascum thapsus*, *Verbena brasiliensis*, *Verbena hastata*, *Verbena hybrids*, *Verbena litoralis*, *Verbena officinalis*, *Verbena rigida*, *Verbesina encelioides*, *Veronica agrestis*, *Veronica chamaedrys*, *Veronica hederifolia*, *Veronica officinalis*, *Veronica persica*, *Vicia amoena*, *Vicia faba*, *Vicia hirsuta*, *Vicia lens*, *Vigna angularis*, *Vigna mungo*, *Vigna radiata*, *Vigna unguiculata* subsp. *sesquipedalis*, *Vigna unguiculata* subsp. *unguiculata*, *Vigna unguiculata*, *Vinca minor*, *Viola cornuta*, *Viola sororia*, *Viola x wittrockiana*, *Vitis vinifera*, *Wahlenbergia marginata*, *Weigela florida*, *Xanthium orientale* subsp. *saccharatum*, *Xanthium spinosum*, *Xanthium strumarium*, *Xerochrysum bracteatum*, *Youngia japonica*, *Yucca aloifolia*, *Yucca baccata*, *Zantedeschia aethiopica*, *Zantedeschia albomaculata* subsp. *albomaculata*, *Zantedeschia albomaculata*, *Zantedeschia elliottiana*, *Zantedeschia rehmannii*, *Zinnia elegans*

GEOGRAPHICAL DISTRIBUTION

Although tomato spotted wilt disease was reported from Australia more than 100 years ago, it only spread rapidly in the 1980s due to the global spread of *Frankliniella occidentalis*, one of the major thrips vectors (EFSA, 2012b; Kormelink *et al.*, 2021). It is now present in almost all countries with temperate, tropical and subtropical climates.



EPPO Region: Albania, Algeria, Armenia, Austria, Azerbaijan, Belgium, Bosnia and Herzegovina, Bulgaria,

Croatia, Cyprus, Czech Republic, Finland, France (mainland), Georgia, Germany, Greece (mainland, Kriti), Guernsey, Hungary, Ireland, Israel, Italy (mainland, Sardegna, Sicilia), Jordan, Lithuania, Malta, Moldova, Montenegro, Netherlands, North Macedonia, Poland, Portugal (mainland, Madeira), Romania, Russia (Far East, Southern Russia), Serbia, Slovenia, Spain (mainland, Islas Baleares, Islas Canarias), Sweden, Switzerland, Tunisia, Türkiye, Ukraine, United Kingdom (Channel Islands, England, Scotland)

Africa: Algeria, Burkina Faso, Congo, Democratic republic of the, Cote d'Ivoire, Egypt, Kenya, Libya, Madagascar, Mauritius, Niger, Nigeria, Reunion, Senegal, South Africa, Sudan, Tanzania, Tunisia, Uganda, Zimbabwe

Asia: Afghanistan, China (Beijing, Chongqing, Gansu, Guangdong, Guizhou, Heilongjiang, Hubei, Liaoning, Ningxia, Qinghai, Shaanxi, Shandong, Sichuan, Tianjin, Yunnan), India (Andhra Pradesh, Assam, Haryana, Himachal Pradesh, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Tamil Nadu, Telangana, Uttar Pradesh), Indonesia (Java), Iran, Iraq, Israel, Japan (Hokkaido, Honshu, Ryukyu Archipelago), Jordan, Korea, Republic, Lebanon, Malaysia (West), Nepal, Oman, Pakistan, Saudi Arabia, Sri Lanka, Syria, Taiwan, Thailand

North America: Canada (Alberta, British Columbia, Manitoba, Nova Scotia, Ontario, Québec, Saskatchewan), Mexico, United States of America (Alabama, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming)

Central America and Caribbean: Costa Rica, Dominican Republic, Haiti, Jamaica, Puerto Rico

South America: Argentina, Bolivia, Brazil (Bahia, Goias, Minas Gerais, Parana, Pernambuco, Sao Paulo), Chile, Colombia, Ecuador, Guyana, Paraguay, Suriname, Uruguay, Venezuela

Oceania: Australia (New South Wales, Northern Territory, Queensland, South Australia, Tasmania, Victoria, Western Australia), Cook Islands, New Zealand, Papua New Guinea

BIOLOGY

TSWV exhibits high genetic diversity (Tsompana *et al.*, 2005; Kaye *et al.*, 2011; Batuman *et al.*, 2020; Fontana *et al.*, 2020; Ruark-Seward *et al.*, 2020). However, despite the known diversity of TSWV populations, it can be assumed that all TSWV isolates occurring worldwide are capable of causing similar diseases in plants and that differences in disease symptoms and symptom severity are a function of isolate, host and environment as well as infection time and the age of the plant (EFSA, 2012a). Diverse resistance-breaking isolates of TSWV have been reported from many regions of the world (Lopez *et al.*, 2011; Almási *et al.*, 2020). The ability of the Tsw gene in pepper and the Sw-5 gene in tomato to break resistance to TSWV is attributed to mutations in different viral genes, NSs for Tsw/pepper (Margarita *et al.*, 2007; de Ronde *et al.*, 2019; Almási *et al.*, 2020) and NSm for Sw-5/tomato (Jahn *et al.*, 2000; Hoffmann *et al.*, 2001; Lopez *et al.*, 2011; Olaya *et al.*, 2020).

Orthotospovirus particles are transmitted and spread in natural conditions by thrips - insects of the genera *Frankliniella* and *Thrips* (family Thripidae). Thrips species known to transmit TSWV are *Frankliniella bispinosa*, *F. cephalica*, *F. fusca*, *F. gemina*, *F. occidentalis*, *F. intonsa*, *F. schultzei*, *Thrips setosus* and *T. tabaci* (Rotenberg & Whitfield, 2018). In the EPPO region, *F. occidentalis*, *F. intonsa* and *T. tabaci* are widely distributed (<https://www.cabi.org/isc/datasheet/>; EFSA, 2012a). Most thrips species are very polyphagous. TSWV is transmitted by thrips in a persistent manner. Only individuals which have acquired the virus at the larval stages can transmit it. Virus transmission occurs after a latent period of circulation and multiplication of TSWV in the thrips vector. Once acquired, the virus is transmitted transstadially and thrips remain infectious for life, however there is no evidence of transovarial transmission (Ullman *et al.*, 1993; Wijkamp *et al.*, 1993; Wijkamp & Peters, 1993; Van de Wetering *et al.*, 1996; Gupta *et al.*, 2018; Rotenberg & Whitfield, 2018).

TSWV can be spread through the movement of infected plant tissue used for vegetative propagation (EFSA, 2012a). Whilst the virus can also be transmitted by mechanical inoculation through small wounds on leaves or stems, this mode of transmission is unlikely to happen in the wild (EFSA, 2012a). Although tospoviruses are considered not to be seed-transmitted (Pappu *et al.*, 2009), an as yet unconfirmed first report of seed transmission of a tospovirus (soybean vein necrosis virus) has recently been published by Groves *et al.* (2016).

DETECTION AND IDENTIFICATION

Symptoms

TSWV can cause a variety of symptoms that may vary in the same host species depending on the variety and age as well as the nutritional and environmental conditions of the plant. Symptoms of TSWV infection also vary depending on the developmental stage of the plant at the time of inoculation and on the virus isolate. Most plants respond to TSWV infection with systemic symptoms. Symptoms on the leaves of infected plants include mosaic, mottle, ring spot, and line patterns. Early infections can result in severe stunting, wilting, leaf deformation and necrosis on stems and tips, chlorotic or necrotic spots on leaves, and plant death. Symptoms on fruits usually consist of irregular discolouration, e.g. yellow/orange flecks or, occasionally, rings, or necrotic lesions or rings. Several other orthospoviruses, as well as even more genetically distinct viruses, can cause symptoms similar to those of TSWV infection. However, it should be noted that infection of host plants by TSWV may be asymptomatic in some host plants, or the symptoms may be mild due to environmental conditions, or the infection may be too recent to reach full symptom expression (Kormelink *et al.*, 2005; EFSA, 2012a,b; Batuman *et al.*, 2020).

On tomatoes, plants show bronzing, curling, necrotic streaks and spots on leaves. Dark brown streaks also appear on leaf petioles, stems and growing tips. Infected plants are small and stunted compared to healthy plants. The ripe fruit shows paler red or yellow areas on the skin. Sometimes infected plants die due to severe necrosis. On pepper, symptoms usually consist of stunting and yellowing of the entire plant. Leaves may show chlorotic line patterns or mosaic with necrotic spots. Necrotic streaks appear on the stems extending to the terminal shoots. On ripe fruits, yellow spots with concentric rings or necrotic streaks have been observed. On lettuces, infection begins on one side of the plant in the leaves, which become chlorotic with brown spots. The discolouration extends to the heart leaves, and growth on the affected side of the plant ceases (Cho *et al.*, 1989; Moriones *et al.*, 1998; Saidi & Warade, 2008; Kamberoglu & Alan, 2011; Salem *et al.*, 2012; Sevik & Arli-Sokmen, 2012; Salamon *et al.*, 2016; Abadkhah *et al.*, 2018; Batuman *et al.*, 2020; Fontana *et al.*, 2020; Kil *et al.*, 2020).

On leaves of ornamentals, the most typical symptoms indicative of TSWV infection are concentric chlorotic to necrotic rings or ring patterns. In some ornamentals (e.g., *Phalaenopsis* spp.), symptoms are restricted to a few leaves, while in others (e.g., chrysanthemums) systemic infection with spots and rings on leaves and systemic necrosis is observed. On chrysanthemums, there is a wide variation of symptoms among cultivars. In other ornamentals, black stem streaks and wilting are usually observed (Cho *et al.*, 1989; Verhoeven and Roenhorst, 1994; Baker *et al.*, 2007; Zheng *et al.*, 2008).

Morphology

TSWV virions are spherical, enveloped particles approximately 80-120 nm in diameter (Kormelink, 2005). The lipid envelope of orthospoviruses contains transmembrane tips composed of two glycoproteins that form oligomeric structures on the outside of the envelope (Butkovi? *et al.*, 2021). The cytoplasmic tails of the glycoproteins interact with the nucleoproteins encapsulating at least one copy of the three linear ssRNA segments (Butkovi? *et al.*, 2021).

Detection and inspection methods

The plants, especially the leaves and fruits, should be examined for symptoms. Particular attention should be paid if thrips are present. If necessary, samples should be taken for laboratory testing for definitive identification of the pest. Procedures have been developed for the inspection of potatoes (EPPO, 2007) and vegetable plants for planting (EPPO, 2016).

The tests recommended for the detection and identification of TSWV are described in EPPO Standard PM7/139 (EPPO, 2020). Electron microscopy can be used for the detection of TSWV and other viruses of the same genus, as they share a typical morphology. Mechanical inoculation of test plants can be used for its detection and subsequent identification by other methods. Several ELISA kits and serological tests for on-site detection are commercially available and can be used as screening tools for TSWV. Several conventional and real-time RT-PCR tests have been described for the detection of TSWV, and some for identification. Sequence analysis of amplicons obtained by the generic conventional PCR assays can also be used for TSWV identification. The definite identification should be based on the sequence of the complete N gene according to the species demarcation criteria of ICTV. In addition, high-throughput sequencing is a technology that can obtain (nearly) complete genome sequences, and analysis of

these sequences can be used to identify a virus isolate.

In 2020, 76 tests for TSWV diagnosis were evaluated in the EU VALITEST project (www.valitest.eu) after an extensive search of scientific papers and commercial providers of tests for the detection of plant pathogens. After a thorough literature search, in silico analyses, in-house testing and test performance study, the most suitable tests for the detection and identification of TSWV in symptomatic tomato leaves were identified (Vučković *et al.*, 2021), and they are listed in the [EPPO Diagnostic Expertise Database](#).

PATHWAYS FOR MOVEMENT

TSWV is a systemic pathogen and as such it is transmitted very efficiently by all vegetative propagation techniques. In international trade, TSWV can be transmitted by susceptible host plants for planting (whether potted or not), and is particularly able to spread if these plants also carry vectors. Thrips are easily transported on above-ground fresh plant parts (e.g. cut foliage, cut flowers and cut branches) hidden under bracts, in buds and leaf bases. Insect vectors that have acquired TSWV and that invade new areas can transmit TSWV to new hosts. Short distance spread of thrips vectors by natural means within and between adjacent greenhouses, orchards, and other production sites is likely, but long-distance spread, even in strong winds is less common. Weeds play an important role in the spread and survival of TSWV through the year; they provide a virus reservoir from which thrips vectors can migrate into crop fields, which then become heavily infected. Mechanical transmission of TSWV is inefficient in the wild and has little impact on agriculture, and transmission through seeds is considered not to be a pathway or to be a highly unlikely pathway (see Biology) (Mound, 1983; Kirk and Terry, 2003; Kormelink, 2005; EFSA, 2012a; Macharia *et al.*, 2016; Batuman *et al.*, 2020).

PEST SIGNIFICANCE

Economic impact

TSWV ranks second in the list of the ten most economically important plant viruses (Scholthof *et al.* 2011; Rybicki, 2015). TSWV became a major agricultural pest in the 1980s with global losses estimated at over one billion USD annually (Goldbach and Peters, 1994). The continued economic importance of TSWV results from its global distribution and wide host range, including many food and ornamental crops important to the EPPO region, significant crop losses due to infection, and the difficulty in controlling thrips and thus the virus (Adkins, 2000; Pappu *et al.*, 2009; Scholthof *et al.*, 2011).

Severe yield and quality losses in tomato were reported by Moriones *et al.* (1998) in trials with TSWV infection in experimental fields in northern Spain. Field trials of tomato in Turkey with TSWV infection in experimental fields resulted in yield losses of up to 42 % with almost complete loss of marketable tomato due to unsightly fruit (Sevik & Arli-Sokmen, 2012). Although it is difficult to draw an inference for actual field situations from studies in experimental fields, TSWV is considered a very serious pathogen of tomato, and severe losses in tomato production have been recorded in Italy, Spain, Bulgaria and Greece (EFSA, 2012a). A similar high impact on a number of other crops, such as peppers, potatoes, eggplants, lettuce and broad beans (*Vicia faba*), has been observed elsewhere (EFSA, 2012a). In Hawaii, for example, TSWV destroyed 50-90% of lettuce crops in some years (Cho *et al.*, 1987). In India, TSWV was reported to be the major viral disease of groundnut (*Arachis hypogaea*) - crop losses ranged from 5 to 80% (Ghanekar *et al.*, 1979). In California, high incidence of TSWV was observed in fields of radicchio (*Cichorium intybus*) (up to 90%), lettuce (15-100%) and peppers (>70%) (Batuman *et al.*, 2020). TSWV infections can also have a major impact on the ornamental industry as the virus is commonly found in greenhouse flower crops (Verhoeven and Roenhorst, 1994). Severe symptoms on leaves and stem necrosis affect the quality of potted plants of Pelargonium, Begonia, Impatiens, Streptocarpus and Chrysanthemum, making them unsaleable (Verhoeven and Roenhorst, 1994). For example, the disease incidence of TSWV on chrysanthemum plants was estimated at 40% in Serbia (Stanković *et al.*, 2013), and at 30% in India (Renukadevi *et al.*, 2015).

Control

The use of healthy planting material and a management strategy based on a combination of thrips and weed control, the use of resistant plant varieties and other measures are considered crucial for TSWV control (EFSA, 2012a;

Batuman *et al.*, 2020).

TSWV is included in the certification scheme for potatoes (EPPO, 1999) and for herbaceous ornamentals (EPPO, 2008), among which there are specific certification schemes for chrysanthemums, pelargoniums, begonias, New Guinea hybrids of impatiens, kalanchoe and petunias (EPPO, 2000a-f). This reduces the impact and spread associated with the plants for the planting pathway.

Seedling beds should be separated from flowering ornamental plants or susceptible crops and surrounding areas kept free of weeds. Greenhouse and outdoor crops should be inspected regularly and as often as possible after planting, and can be tested during the entire propagation scheme. The presence of thrips in crops should be monitored with yellow sticky card traps. If the disease appears in a crop, infected plants should be immediately uprooted and destroyed, and the greenhouse/ field treated with an insecticide against thrips (EFSA, 2012a; EPPO, 2008; Batuman *et al.*, 2020).

Thrips are difficult to control because they can become resistant to several insecticides used (Gao *et al.*, 2012). To avoid the development of resistance mechanisms, it is important to rotate insecticides with different active ingredients (Bielza, 2008). The prevalence of thrips vectors in some crops (e.g. pepper and tomato) under protected conditions can be reduced by introducing biocontrol agents. These include predatory mites, entomopathogenic fungi and nematodes, parasitic wasps and generalist predators (Sánchez and Lacasa, 2002; Blaaser *et al.*, 2004; Ansari *et al.*, 2008; Messelink *et al.*, 2008; Cloyd, 2009). To prevent infestation of a new crop with a thrips population, it is recommended that greenhouses be kept completely free of crop residues and weeds for at least six weeks prior to planting and that seedlings be treated with insecticides before planting (EFSA, 2012a). In addition, fine-mesh netting can be potentially useful to exclude thrips in greenhouses (EFSA, 2012a).

Cultivation of TSWV resistant (tolerant/immune) cultivars will reduce the incidence of the disease (Pico *et al.*, 2002; Saidi and Warade, 2008; Dianese *et al.*, 2011; de Oliveira *et al.*, 2018; Padmanabhan *et al.*, 2019). Unfortunately, resistant varieties of the vast majority of TSWV host plants are not available (EFSA, 2012a), and there is a risk of selection for virulent resistance-breaking strains, as in the case of the dominant gene Sw-5 for TSWV resistance in tomato and the gene Tsw for TSWV resistance in pepper (see Biology).

Phytosanitary risk

TSWV is reported in many European and Mediterranean countries and has one of the largest known host ranges of any plant virus, including cultivated and wild plants. In addition, its thrips vector species are widely distributed in these countries. Spread can occur through thrips vector activity and through movement of infected hosts. The only major limiting factors are climatic conditions in Northern Europe which appear to limit the capacity of thrips vectors to develop and persist in the open environment. Since TSWV can cause both yield and quality losses in a wide range of hosts, without appropriate control measures, significant impact is expected, under protected conditions, or in open fields in Southern Europe (EFSA, 2012a).

PHYTOSANITARY MEASURES

To prevent the introduction and spread of TSWV, most countries apply import requirements to some host species. These requirements vary depending on the crop and the virus prevalence at the point of origin. Following deregulation as an EU quarantine pest, TSWV was recommended for regulation as a regulated non-quarantine pest for seed potatoes, propagating and planting material (other than seeds) of some vegetables (pepper, lettuce, tomato and eggplant), for tobacco and for some ornamentals (*Begonia x hiemalis*, *Gerbera*, Pelargonium, *Capsicum annuum*, *Chrysanthemum* and Impatiens New Guinea hybrids) (Picard *et al.*, 2018).

EPPO countries where TSWV does not occur or is not widespread may consider regulation. If they do so, these countries may require that plants have been produced in a pest free area, in a pest-free place/site of production or be subject to inspection and/or testing (EFSA, 2012a) e.g. as performed in certification schemes. In particular, EPPO recommends that seed potatoes to be imported should meet the requirements of EPPO Standard PM 4/28 *Certification scheme for seed potatoes* (or equivalent) (EPPO, 1999) and be shown free from insects by visual inspection (EPPO, 2017). Post-entry quarantine programmes are established to allow the safe movement of germplasm for research and breeding purposes. During post-entry quarantine for potatoes, it is recommended that

testing for viruses be carried out on microplants and/or greenhouse-grown plants; in exceptional cases, tubers (tuber sap or sprouts) may also be tested (EPPO, 2019).

Before starting the propagation and/or production of plants, it should be ensured that the planting material is free from TSWV (TSWV is included in certification schemes for potatoes and various ornamental plants, see Control section). During the EU Quality pest project, in addition to the testing of nuclear stock (or the stipulation that it is derived from tested mother plants), recommended measures for all other plants for planting categories consisted of zero-tolerance based on symptom and/or testing, and active control of the vectors at the place of production.

REFERENCES

Abadkhah M, Koolivand D & Eini O (2018) A new distinct cClade for Iranian Tomato spotted wilt virus isolates based on the polymerase, nucleocapsid, and non-structural genes. *The Plant Pathology Journal* **34**(6), 514-531.

Abudurexiti A, Adkins S, Alioto D Alkhovsky SV, Avšič-Županc T, Ballinger MJ, Bente DA, Kuhn JH *et al.* (2019) Taxonomy of the order Bunyavirales: update 2019. *Archives of Virology*. <https://doi.org/10.1007/s00705-019-04253-6>

Adkins S (2000) Tomato spotted wilt virus—positive steps to negative success. *Molecular Plant Pathology* **1**, 151–157.

Almási A, Nemes K & Salánki K (2020) Increasing diversity of resistance breaking pepper strains of Tomato spotted wilt virus in the Mediterranean region. *Phytopathologia Mediterranea* **59**(2), 385-391.

Ansari M, Brownbridge M, Shah F & Butt T (2008) Efficacy of entomopathogenic fungi against soil-dwelling life stages of western flower thrips, *Frankliniella occidentalis*, in plant-growing media. *Entomologia Experimentalis et Applicata* **127**, 80–87.

Baker CA, Davison D & Jones L (2007) Impatiens necrotic spot virus and Tomato spotted wilt virus diagnosed in *Phalaenopsis* orchids from two Florida nurseries. *Plant Disease* **91**, 1515.

Batuman O, Turini TA, LeStrange M, Stoddard S, Miyao G, Aegerter BJ, Chen L-F, McRoberts N, Ullman DE & Gilbertson RL (2020) Development of an IPM strategy for Thrips and Tomato spotted wilt virus in processing tomatoes in the Central Valley of California. *Pathogens* **9**, 636. <https://doi.org/10.3390/pathogens9080636>

Bielza P (2008) Insecticide resistance management strategies against the western flower thrips, *Frankliniella occidentalis*. *Pest Management Science* **64**, 1131–1138.

Blaeser P, Sengonca C & Zegula T (2004) The potential use of different predatory bug species in the biological control of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae). *Journal of Pest Science* **77**, 211–219.

Butkovič A, González R & Elena SF (2021) Revisiting Orthotospovirus phylogeny using full-genome data and testing the contribution of selection, recombination and segment reassortment in the origin of members of new species. *Archives of Virology* **166**, 491–499.

Cho JJ, Mitchell WC, Mau RFL & Sakimura K (1987) Epidemiology of tomato spotted wilt virus disease on crisphead lettuces in Hawaii. *Plant Disease* **71**, 505-508.

Cho JJ, Mau RFL, German TL, Hartmann RW, Yudin LS, Gonsalves D & Provvidenti R (1989) A multidisciplinary approach to management of tomato spotted wilt virus in Hawaii. *Plant Disease* **73**(5), 375-383.

Cloyd RA (2009) Western flower thrips (*Frankliniella occidentalis*) management on ornamental crops grown in greenhouses: Have we reached an impasse? *Pest Technology* **3**, 1–9.

de Oliveira AS, Boiteux LS, Kormelink R & Resende RO (2018) The Sw-5 Gene Cluster: Tomato breeding and research toward *Orthotospovirus* disease control. *Frontiers in Plant Science* **9**, 1055. <https://doi.org/10.3389/fpls.2018.01055>

- de Ronde D, Lohuis D & Kormelink R (2019) Identification and characterization of a new class of Tomato spotted wilt virus isolates that break Tsw-based resistance in a temperature-dependent manner. *Plant Pathology* **68**, 60-71.
- Dianese EC, Fonseca MEN, Inoue-Nagata AK, Resende RO & Boiteux LS (2011) Search in *Solanum* (section *Lycopersicon*) germplasm for sources of broad-spectrum resistance to four *Tospovirus* species. *Euphytica* **180**, 307–319.
- EFSA Panel on Plant Health (2012a) Scientific Opinion on the risk to plant health posed by Tomato spotted wilt virus to the EU territory with identification and evaluation of risk reduction options. *EFSA Journal* **10**(12), 3029. <https://doi.org/10.2903/j.efsa.2012.3029>
- EFSA Panel on Plant Health (2012b) Scientific Opinion on the pest categorisation of the tospoviruses. *EFSA Journal* **10**(7), 2772. <https://doi.org/10.2903/j.efsa.2012.2772>
- EPPO (1999) Certification schemes. PM 4/28 (1) Seed potatoes. *EPPO Bulletin* **29**, 253-267.
- EPPO (2000a) Production of healthy plants for planting. PM 4/6 (2) Certification scheme for chrysanthemum. *EPPO Bulletin* **32**, 105-114.
- EPPO (2000b) Production of healthy plants for planting. PM 4/3 (3) Certification scheme for pelargonium. *EPPO Bulletin* **32**, 67-78.
- EPPO (2000c) Production of healthy plants for planting. PM 4/19 (2) Certification scheme for begonia. *EPPO Bulletin* **32**, 135-145.
- EPPO (2000d) Production of healthy plants for planting. PM 4/20 (2) Certification scheme for New Guinea hybrids of impatiens. *EPPO Bulletin* **32**, 147-157.
- EPPO (2000e) Production of healthy plants for planting. PM 4/25 (2) Certification scheme for kalanchoe. *EPPO Bulletin* **32**, 199-210.
- EPPO (2000f) Production of healthy plants for planting. PM 4/26 (2) Certification scheme for petunia. *EPPO Bulletin* **32**, 211-221.
- EPPO (2007) Phytosanitary procedures. PM 3/71 General crop inspection procedure for potatoes. *EPPO Bulletin* **37**, 592-597.
- EPPO (2008) Schemes for the production of healthy plants for planting. PM 4/34 (1) Production of pathogen-tested herbaceous ornamentals. *EPPO Bulletin* **38**, 31-52.
- EPPO (2016) Phytosanitary procedures. PM 3/77 (1) Vegetable plants for planting under protected conditions – inspection of places of production. *EPPO Bulletin* **46**, 40-48.
- EPPO (2017) Commodity-specific phytosanitary measures. PM 8/1 (2) Potato. *EPPO Bulletin* **47**, 487-503.
- EPPO (2019) Phytosanitary procedures. PM 3/21 (3) Post entry quarantine for potato. *EPPO Bulletin* **49**, 452-479.
- EPPO (2020) PM 7/139(1) Tospoviruses (Genus *Orthotospovirus*). *EPPO Bulletin* **50**, 217-240.
- Fontana A, Albanese G, Mangalli A, Tomassoli L & Tiberini A (2020) Phylogenetic analysis based on full genome sequencing of Italian tomato spotted wilt virus isolates identified in “Roggianese” sweet pepper and chilli pepper. *Annals of Applied Biology* **176**, 170-179.
- Gao Y, Lei Z & Reitz SR (2012) Western flower thrips resistance to insecticides: detection, mechanisms and management strategies. *Pest Management Science* **68**, 111-1121.
- Ghanekar AM, Reddy DVR, Iizuka N, Amin PW & Gibbons RW (1979) Bud necrosis of groundnut (*Arachis hypogaea*)

) in India caused by tomato spotted wilt virus. *Annals of Applied Biology* **93**, 173-179.

Goldbach R & Peters D (1994) Possible causes of the emergence of tospovirus diseases. *Seminars in Virology* **5**, 113-120.

Groves C, German T, Dasgupta R, Mueller D & Smith DL (2016) Seed transmission of Soybean vein necrosis virus: The first Tospovirus implicated in seed transmission. *PLoS One* **11**(1), e0147342.
<https://doi.org/10.1371/journal.pone.0147342>

Gupta R, Kwon S-Y & Kim ST (2018) An insight into the tomato spotted wilt virus (TSWV), tomato and thrips interaction. *Plant Biotechnology Reports* **12**, 157-163.

Hoffmann K, Qiu WP & Moyer JW (2001) Overcoming host- and pathogen-mediated resistance in tomato and tobacco maps to the M RNA of Tomato spotted wilt virus. *Molecular Plant-Microbe Interactions* **14**, 242-249.

Jahn M, Paran I, Hoffmann K, Radwanski ER, Livingstone KD, Grube RC, Aftergoot E, Lapidot M & Moyer J, (2000) Genetic mapping of the Tsw locus for resistance to the Tospovirus Tomato spotted wilt virus in *Capsicum* spp. and its relationship to the Sw-5 gene for resistance to the same pathogen in tomato. *Molecular Plant-Microbe Interactions* **13**(3), 673-682.

Kamberoglu MA & Alan B (2011) Occurrence of Tomato spotted wilt virus in lettuce in cukurova region of Turkey. *International Journal of Agriculture & Biology* **13**, 431-434.

Kaye AC, Moyer JW, Parks EJ, Carbone I & Cubeta MA (2011) Population genetic analysis of Tomato spotted wilt virus on peanut in North Carolina and Virginia. *Phytopathology* **101**, 147- 153.

Kil EJ, Chung YJ, Choi HS, Lee S & Kim CS (2020) Life Cycle-Based Host Range Analysis for Tomato Spotted Wilt Virus in Korea. *The Plant Pathology Journal* **36**(1), 67-75.

Kirk WDJ & Terry LI (2003) The spread of the western flower thrips *Frankliniella occidentalis* (Pergande). *Agricultural and Forest Entomology* **5**, 301-310.

Kormelink R (2005) Tomato spotted wilt virus. In: Description of plant viruses (DPV) no. 412, <https://dpvweb.net/> (accessed on 5 June 2021)

Kormelink R, Verchot J, Tao X & Desbiez C (2021) The Bunyavirales: The Plant-Infecting Counterparts. *Viruses* **13**, 842, <https://doi.org/10.3390/v13050842>

Lopez C, Aramburu J, Galipienso L, Soler S, Nuez F & Rubio L (2011) Evolutionary analysis of tomato Sw-5 resistance-breaking isolates of Tomato spotted wilt virus. *Journal of General Virology* **92**, 210-215.

Macharia I, Backhouse D, Wu S-B & Ateka EM (2016) Weed species in tomato production and their role as alternate hosts of Tomato spotted wilt virus and its vector *Frankliniella occidentalis*. *Annals of Applied Biology* **169**, 224-235.

Margaria P, Ciuffo M, Pacifico D & Turina M (2007) Evidence that the nonstructural protein of Tomato spotted wilt virus is the avirulence determinant in the interaction with resistant pepper carrying the TSW gene. *Molecular Plant-Microbe Interactions* **20**, 547-558.

Messelink GJ, Maanen R, van Steenpaal SEF & Janssen A (2008) Biological control of thrips and whiteflies by a shared predator: two pests are better than one. *Biological Control* **44**, 372-379.

Moriones E, Aramburu J, Riudavets J, Arno J & Lavina A (1998) Effect of plant age at time of infection by tomato spotted wilt tospovirus on the yield of field-grown tomato. *European Journal of Plant Pathology* **104**, 295-300.

Mound LA (1983) Natural and disrupted patterns of geographical distribution in Thysanoptera (Insecta). *Journal of Biogeography* **10**(2), 119-133.

- Olaya C, Fletcher SJ, Zhai Y, Peters J, Margaria P, Winter S, Mitter N & Pappu HR (2020) The Tomato spotted wilt virus (TSWV) genome is differentially targeted in TSWV-infected tomato (*Solanum lycopersicum*) with or without Sw-5 Gene. *Viruses* **12**, 363. <https://doi.org/10.3390/v12040363>
- Padmanabhan C, Ma Q, Shekasteband R, Stewart KS, Hutton SF, Scott JW, Fei Z & Ling KS (2019) Comprehensive transcriptome analysis and functional characterization of PR-5 for its involvement in tomato Sw-7 resistance to tomato spotted wilt tospovirus. *Scientific reports* **9**, 7673. <https://doi.org/10.1038/s41598-019-44100-x>
- Pappu HR, Jones RAC & Jain RK (2009) Global status of tospovirus epidemics in diverse cropping systems: successes achieved and challenges ahead. *Virus Research* **141**, 219–236.
- Parrella G, Gognalons P, Gebre-Selassie K, Vovlas C & Marchoux G (2003) An update of the host range of tomato spotted wilt virus. *Journal of Plant Pathology* **85**(4), 227-264.
- Picard D, Afonso T, Benko-Beloglavec A, Karadjova O, Matthews-Berry S, Paunovic SA, Pietsch M, Reed P, van der Gaag DJ & Ward M (2018) Recommended regulated non-quarantine pests (RNQPs), associated thresholds and risk management measures in the European and Mediterranean region. *EPPO Bulletin* **48**, 552-558.
- Pico B, Herraiz J, Ruiz JJ & Nuez F (2002) Widening the genetic basis of virus resistance in tomato. *Scientia Horticulturae* **94**, 73–89.
- Plyusnin AMQ, Beaty BJ, Elliott RM, Goldbach R, Kormelink R, Lundkvist A, Schmaljohn CS & Tesh RB (2011) Bunyaviridae. In: *Virus taxonomy, ninth report of the International Committee on Taxonomy of Viruses*. Elsevier Academic Press (Eds.: King MJA, Adams MJ, Carstens EB & Lefkowitz EJ) London, UK, 725–741.
- Renukadevi P, Nagendran K, Nakkeeran S, Karthikeyan G, Jawaharlal M, Alice D, Malathi VG & Pappu HR (2015) First Report of Tomato spotted wilt virus Infection of Chrysanthemum in India. *Plant Disease* **99**(8), 1190.
- Rotenberg D & Whitfield AE (2018) Molecular interactions between tospoviruses and thrips vectors. *Current Opinion in Virology* **33**, 191–197.
- Ruark?Seward C, Bonville B, Kennedy G & Rasmussen DA (2020) Evolutionary dynamics of Tomato spotted wilt virus within and between alternate plant hosts and thrips. *Scientific Reports* **10**, 15797. <https://doi.org/10.1038/s41598-020-72691-3>
- Rybicki EP (2015) A Top Ten list for economically important plant viruses. *Archives of Virology* **160**, 17–20.
- Salem NM, Mansour A & Badwan H (2012) Identification and partial characterization of tomato spotted wilt virus on lettuce in Jordan. *Journal of Plant Pathology* **94**(2), 431-435.
- Samuel G, Bald JG & Pittman HA (1930) Investigations on 'spotted wilt' of tomatoes. *Australian Council of Science and Industrial Research Bulletin* **44**, 64 pp.
- Saidi M & Warade S (2008) Tomato breeding for resistance to Tomato spotted wilt virus (TSWV): an overview of conventional and molecular approaches. *Czech Journal of Genetics and Plant Breeding* **44**, 83–92.
- Salamon P, Mityko J, Kalo P & Szabo Z (2016) Symptoms caused by Tomato spotted wilt virus (TSWV) in pepper (*Capsicum* spp.) and marker assisted selection of TSWV resistant pepper lines for hybrid constructions. *Proceedings XVI. EUCARPIA Capsicum and Eggplant Meeting*, Kecskemét, Hungary, 12-14. Sept. 2016. pp. 69-75.
- Sánchez JA & Lacasa A (2002) Modelling population dynamics of *Orius laevigatus* and *O. albidipennis* (Hemiptera: Anthocoridae) to optimise their use as biological control agents of *Frankliniella occidentalis* (Thysanoptera: Thripidae). *Bulletin of Entomological Research* **92**, 77– 78.
- Scholthof KB, Adkins S, Czosnek H, Palukaitis P, Jacquot E, Hohn T, Hohn B, Saunders K, Candresse T, Ahlquist P, Hemenway C & Foster GD (2011) Top 10 plant viruses in molecular plant pathology. *Molecular Plant Pathology* **12**, 938–954.

- Sevik MA & Arli-Sokmen M (2012) Estimation of the effect of Tomato spotted wilt virus (TSWV) infection on some yield components of tomato. *Phytoparasitica* **40**, 87–93.
- Stankovi? I, Bulaji? A, Vu?urovi? A, Risti? D, Milojevi? K, Nikoli? D & Krsti? B (2013) First report of Tomato spotted wilt virus on Chrysanthemum in Serbia. *Plant Disease* **97**, 150.
- Tsompana M, Abad J, Purugganan M & Moyer JW (2005) The molecular population genetics of the Tomato spotted wilt virus (TSWV) genome. *Molecular Ecology* **14**, 53–66.
- Ullman DE, German TL, Sherwood JL, Westcot DM & Cantone FA (1993) Tospovirus replication in insect vector cells: Immunocytochemical evidence that the nonstructural protein encoded by the S RNA of tomato spotted wilt tospovirus is present in thrips vector cells. *Phytopathology* **83**, 456–463.
- van de Wetering F, Goldbach R & Peters D (1996) Tomato spotted wilt tospovirus ingestion by first instar larvae of *Frankliniella occidentalis* is a prerequisite for transmission. *Phytopathology* **86**, 900–905.
- Verhoeven TJ & Roenhorst JW (1994) Tomato spotted wilt virus: ecological aspects in ornamental crops in the Netherlands from 1989 up to 1991. *Acta Horticulturae* **377**, 175–182.
- Vu?urovi? A, Jakoš N, Ravnikar M & Mehle N (2021) Results of the test performance study on the detection and identification of tomato spotted wilt tospovirus: re-emerging threat to vegetables and ornamentals. In: *Programme and abstracts: International Advances in Plant Virology 2021*, virtually via Zoom, 20th April - 22nd April 2021. *Association of Applied Biologists*, 23-24.
- Wijkamp I & Peters D (1993) Determination of the median latent period of 2 Tospoviruses in *Frankliniella occidentalis*, using a novel leaf disk assay. *Phytopathology* **83**, 986–991.
- Wijkamp I, van Lent J, Kormelink R, Goldbach R & Peters D (1993) Multiplication of tomato spotted wilt virus in its insect vector, *Frankliniella occidentalis*. *The Journal of general virology* **74**, 341.
- Zheng YX, Chen CC, Yang CJ, Yeh SD & Jan FJ (2008) Identification and characterization of a tospovirus causing chlorotic ringspots on *Phalaenopsis* orchids. *European Journal of Plant Pathology* **120**, 199–209.

CABI resources used when preparing this datasheet

CABI Datasheet on *Frankliniella occidentalis* (western flower thrips) (<https://www.cabi.org/isc/datasheet/24426>; date of the last modification: 22 September 2020; accessed on June 2021)

ACKNOWLEDGEMENTS

This datasheet was extensively revised in 2021 by Nataša Mehle (National Institute of Biology, Slovenia). Her valuable contribution is gratefully acknowledged.

How to cite this datasheet?

EPPO (2024) *Orthotospovirus tomatomaculae*. EPPO datasheets on pests recommended for regulation. Available online. <https://gd.eppo.int>

Datasheet history

This datasheet was first published in 1992 and revised in 1997 and 1999, as well as in 2021. It is now maintained in an electronic format in the EPPO Global Database. The sections on 'Identity', 'Hosts', and 'Geographical distribution' are automatically updated from the database. For other sections, the date of last revision is indicated on the right.

CABI/EPPO (1992/1997) *Quarantine Pests for Europe (1st and 2nd edition)*. CABI, Wallingford (GB).

EPPO (1999) EPPO Data sheets on quarantine pests - Tomato spotted wilt tospovirus. *EPPO Bulletin* **29**(4), 465-472.



Co-funded by the
European Union