

Report of a Pest Risk Assessment

This summary presents the main features of a pest risk assessment which has been conducted on the pest, according to EPPO Standard PP 5/3(1) Pest Risk Assessment Scheme.

Pest: *Dasychira albodentata*
PRA area: The European part of the EPPO region
Assessor: EPPO Secretariat
Date: June, 2000

1. INITIATION

1.1 Reason for doing PRA: Study of the risk of forest pests occurring on the territory of the former USSR for the western part of EPPO region
1.2. Taxonomic position of pest: *Dasychira albodentata* Bremer (Lepidoptera: Lymantriidae)

2. PROBABILITY OF INTRODUCTION

2.1 Entry

2.1.1 Geographical distribution: Of limited distribution in EPPO region
Originates in Russia (Transbaikalia and surrounding regions)
Europe: Absent
Asia: Russia (east of Southern Siberia, Transbaikalia, south of North – Eastern Siberia, Southern Far East), China (north), Mongolia (north)
North America: Absent
Central America & Caribbean: Absent
South America: Absent
Oceania: Absent

2.1.2 Major host plants: Several species of *Pinus*, *Larix* and some other coniferous. Its preferred hosts are *Larix gmelinii* (= *Larix dahurica*), *Pinus sylvestris*, *Pinus pumila* and *Pinus koraiensis*.

2.1.3 Which pathway(s) is the pest likely to be introduced on: All stages of the life cycle of *D. albodentata* can be transported on plants moving in trade particularly plants for planting and cut

branches (including Christmas trees). Eggs and larvae may be associated with wood containing bark and may be hitchhikers on other products.

In decreasing order of risk, pathways for *D. albodentata* may be:

1. Untreated wood with bark
2. Host plants for planting and cut branches
3. Wood without bark, dunnage and packing material
4. Ships, planes, trains, road transports

2.2 Establishment

2.2.1 Crops at risk in the PRA area:

All species of *Pinus*, *Larix* and other coniferous trees. The biggest risk exists for forests.

2.2.2 Climatic similarity of present distribution with PRA area (or parts thereof):

East, north and centre of the European part of the EPPO region has a similar climatic conditions with the area of origin and present distribution of the pest

2.2.3 Aspects of the pest's biology that would favour establishment:

The pest is polyphagous and genetically adaptable

2.2.4 Characteristics (other than climatic) of the PRA area that would favour establishment:

Host plants are widely distributed within the PRA area. Suitable ecological niches are available throughout the PRA area.

2.2.5 Which part of the PRA area is the endangered area:

The endangered part of the PRA area covers primarily eastern, northern and central parts of the European EPPO region (Austria, Belgium, Czech Republic, Estonia, Finland, France, Germany, Hungary, Latvia, Lithuania, Norway, Poland, Slovakia, Sweden, Switzerland) as well as mountain areas of some other countries.

3. ECONOMIC IMPACT ASSESSMENT

3.1 Describe damage to potential hosts in PRA area:

D. albodentata attacks both stressed and healthy trees of different ages leading to decrease of wood and seed production and sometimes their death and/or to outbreaks of bark beetles and other pests.

3.2 How much economic impact does the pest have in its present distribution:

D. albodentata is an important defoliator of coniferous trees in the region of its present distribution. Its outbreaks occur throughout large areas (thousands of hectares), cause decrease of wood and seed production and sometimes lead to the death of forests. The reforestation of these areas is complicated and takes much time. This results in serious changes of environment over large areas. The death of forests has a social influence on the people living in damaged areas. Large scale pesticide treatments influence the social value of forest berries and mushrooms.

3.3 How much economic impact would the pest have in the PRA area:

Considering the similarity of ecological conditions, the damage in the PRA area should be not much less than in the present area of the pest.

4. CONCLUSIONS OF PRA

4.1 Summarize the major factors that influence the acceptability of the risk from this pest:

This pest

- comes from an area with similar climatic conditions to the PRA area and causes serious economic damage there;
- could easily establish throughout a part of PRA area;
- is the pest of coniferous trees which are very important in the PRA area;
- can cause also environmental and social damage.

4.2 Estimate the probability of entry:

medium (from 3.91 to 5.45 for different pathways)

4.3 Estimate the probability of establishment:

high (6.57)

4.4 Estimate the potential economic impact:

medium (4.89)

4.5 Degree of uncertainty

There is little uncertainty in this assessment

5. OVERALL CONCLUSIONS OF THE ASSESSOR

The endangered area is primarily eastern, northern and central parts of the European EPPO region (Austria, Belgium, Czech Republic, Estonia, Finland, France, Germany, Hungary, Latvia, Lithuania, Norway, Poland, Slovakia, Sweden, Switzerland) as well as mountain areas of some other countries. The potential impact within the endangered area is rather high including both the direct damage to coniferous plantations and forests (mainly *Pinus* and *Larix*) resulting in wood and seed losses, environmental damage to natural forests resulting in their death on large areas, and social damage to people living in damaged areas.

D. albobdentata should be included into the A2 EPPO list.