

European and Mediterranean Plant Protection Organization
 Organisation Européenne et Méditerranéenne pour la Protection des Plantes

Data sheets on pests recommended for regulation

Fiches informatives sur les organismes recommandés pour réglementation

Ambrosia confertiflora DC

Identity

Scientific name: *Ambrosia confertiflora* DC.

Synonyms: *Ambrosia caudata* (Rydb.) Shinnery, *Ambrosia simulans* Shinnery, *Franseria caudata* Rydb., *Franseria confertiflora* (DC.) Rydb., *Franseria hispidissima* Rydb., *Franseria incana* Rydb., *Franseria pringlei* Rydb., *Franseria strigulosa* Rydb., *Franseria tenuifolia* Harv. & A.Gray, *Franseria tenuifolia* var. *tripinnatifida* A.Gray, *Gaertneria tenuifolia* (Harv. & A.Gray) Kuntze, *Gaertneria tenuifolia* Harv. & A.Gray

Taxonomic position: phylum Spermatophyta; subphylum Angiospermae; class Dicotyledonae; order Asterales; family Asteraceae

Common names: English: burr ragweed, slimleaf bursage, weak leaf burr ragweed; French: herbe blanche, Spanish: ajenjo del campo, yerba de la oveja, altamisa falsa; Mexican Spanish: escoba amarga

EPPO Code: FRSCO

Phytosanitary categorization: EPPO A2 List no. 420

Geographical distribution

Native range

North America: Mexico (Aguascalientes, Baja California Norte, Baja California Sur, Chihuahua, Coahuila, Colima, Durango, Guanajuato, Guerrero, Jalisco, México, Nuevo León, Querétaro, San Luis Potosí, Sinaloa, Sonora, Tamaulipas and Zacatecas), USA (Arizona, California, Colorado, Kansas, Nevada, New Mexico, Oklahoma, Tennessee, Texas, Utah)

Caribbean: Puerto Rico

Introduced range

EPPO region: Israel and the West Bank

Oceania: Australia (New South Wales, Queensland, South Australia)

History of introduction and spread

Ambrosia confertiflora is established and invasive in Israel (Dufour-Dror, 2012; see also Dufour-Dror & Yaacoby, 2013; Dufour-Dror, 2016; Dufour-Dror & Fragman-Sapir, 2018). *Ambrosia confertiflora* was first recorded in Israel near to the town of Zichron Yakov in 1990. *Ambrosia confertiflora* began to spread in the Shechem (Nablus) area in central Samaria during the mid-1990s. It is now widespread in Samaria (the northern part of the West Bank), especially along roads and in cultivated areas, but also in nature reserves such as Wadi Qana. The plant spread westwards and is now present in the Heffer Valley area (Sharon region), where 480 ha of riverbanks along the Alexander River are already heavily infested. The plant also spread eastward along Wadi Tirza and has reached the Northern Jordan Valley. Additional foci were recently discovered along the Yarkon River, on the Carmel Mount, in the Haifa region, in the southern part of the Yizre'el Valley in Lower Galilee, in the south near Gadera and in the Ashdod area (Israel Ministry of Environmental Protection, 2013).

Between the late 1990s and the early 2000s, a massive invasion began along the bank of the Alexander River, apparently originating from the area of Sechem (Nablus) in Central Samaria (West Bank). During the 2000s, a heavy infestation was recorded along the Wadi Tirza, flowing from the Sechem area to the east in the Jordan Valley. Since then new occurrences have been recorded along the Jordan Valley, in Lower Galilee (few sites) and in 2017 in the Negev Desert, south of Beer-Sheva.

In Australia, the species was introduced into Queensland in 1950 and is restricted to small colonies in the Western Darling Downs and the Burnett Pastoral District. In New South Wales, populations occur on the Far Western and North-Western Plains and on the Central Western Slopes, while in South Australia the species is only recorded in the surroundings of Adelaide (Parsons & Cuthbertson, 2001). There are very few records, and since Parsons & Cuthbertson (2001) the plant has been mentioned by Richardson *et al.*

(2011), but apparently it does not spread vigorously in this region.

Morphology

Plant type

Erect perennial herb (hemicytrophite).

Description

Ambrosia confertiflora is a perennial species 20–80 (–150+) cm in height. Stems erect leaves opposite at the base, alternating higher up with petioles 10–35 mm; lanceolate to ovate in outline, 40–85(–150) × 20–35(–55) mm overall, lacinate 2–4-pinnately lobed (lobes ± lanceolate); bases cuneate to truncate, ultimate margins entire, abaxial and adaxial faces strigillose to sericeous (often grayish) and gland-dotted. Pistillate heads clustered, proximal to staminate; florets 1(–2). Staminate heads: peduncles 0.5–2 mm; involucre cup-shaped, 1.5–3+ mm in diameter, strigillose; florets 5–20+. Fruit a spiny bur, pyramidal to pyriform, 1–2 mm long, strigillose to pilosulous, spines (1–)5–12+, mostly distal, stoutly conic to subulate, 0.5–1 mm, tips unciniate. $2n = 72, 108$ (Flora of North America, 2017).

The leaves have short petioles and are bipinnate, divided into long lobes; their margins have sparse short hairs. The leaves are opposite at the base of the stem and alternate on the upper part of the plant. The leaves release a characteristic strong smell that may help to identify the plant. *Ambrosia confertiflora* is monoecious and has male and female yellow or greenish flowers. The male flowers are numerous and small (diameter about 1 cm) and are borne on erect clusters (Fig. 1). The female flowers lack petals and are concentrated in the leaf axils in a cup-shaped involucre. The fruit is a burr with 10–20 hooked spines and includes a single seed. The seeds are brown with a diameter of 3–4 mm. In Israel, the species blooms from September through to December.

Biology and ecology

General

Germination is best when seeds are on the soil surface (no germination was detected when seeds were planted at depths below 4 cm) and at moderate temperatures (Yair, 2017), suggesting autumn or spring germination. This is supported by Australian observations that this species germinates mostly in autumn (Parsons & Cuthbertson, 2001). It is recorded that in California (USA) *A. confertiflora* can tolerate a wide pH from 5.4 to 7.4 (Calflora, 2017).

Habitats

In the native range (Arizona), *Ambrosia confertiflora* grows in dry or moist, rocky, sterile or fertile soils. It is abundant



Fig. 1 Flowering *Ambrosia confertiflora*. Photo courtesy of J. M. Dufour.

along city streets, highways, waste places and edges of cultivated fields, also on barren mesas and slopes in southern and central Arizona; it is less troublesome northward, except in local areas where colonies have become established (e.g. around Flagstaff); 300–2000 m elevation.

In Israel, *Ambrosia confertiflora* occurs in various natural and disturbed habitats, including dry plains and semi-arid valleys, degraded pastures, cultivated orchards, summer field crops (cotton and water melon), avocado and date groves, along roadsides, river banks and wadi beds (dry river beds), in wastelands and other disturbed areas (J. M. Dufour, pers. comm. 2017).

In Australia, the species is a weed in degraded pasture land and agricultural land, also occurring in open disturbed areas (Parsons & Cuthbertson, 2001).

Environmental requirements

The establishment of populations in the Jordan Valley, along roadsides, near Jericho and more recently in the Negev suggests that the plant can cope easily with arid conditions, i.e. high summer temperatures and an extended dry season. This is supported by the distribution of this species in its native range where it is most common in arid to semi-arid climates (Calflora, 2017; Flora of North America, 2017). However, it appears to have a wide environmental tolerance in its native range, occurring in temperatures from –1 to 35°C, precipitation levels from 230 to 1040 mm, elevations from 10 to 2000 m in regions where the growing season ranges from 5 to 12 months (Calflora, 2017; Flora of North America, 2017). In Israel, it is commonly associated with open habitats along rivers or in seasonally moist

habitats (Yair, 2017). Growth trials have shown that the plant grows better in warm conditions (22/28°C and 28/34°C, night/day cycle) and shoot elongation is slower in cold conditions (10/16°C and 16/22°C, night/day cycle), and in these conditions the plant remains in a rosette form (Yair, 2017).

Natural enemies

The lepidopteran *Epiblema strenuana* is present in Israel but shows no significant impact on *A. confertiflora* (Yaacoby & Seplyarsky, 2011).

Uses and benefits

To date, within the EPPO region, there are no known socio-economic benefits derived from the plant species *A. confertiflora*. The species is not known to be available via the horticulture industry.

Pathways for movement

There is no clear information on how the species was introduced into Israel. It is also unknown how the species entered Australia. To date the species has not been intercepted along any pathway into the EPPO region. When compiling the pest risk analysis (PRA), the Expert Working Group considered the pathways listed below as the *most feasible* pathways for the entry of the species into the region based on experience with similar non-native plant occurrences (EPPO, 2018):

- shipments of cattle and sheep from Australia to the EPPO region – particularly Israel and Jordan
- seed may become incorporated into animal seed mixture and imported and dispersed within the PRA area. The likelihood of contamination in animal feed is elevated as regulations are less restrictive than for human consumption or seed for planting
- in Israel, this species was recorded in the Golan Heights for the first time at a development site where the machinery is known to have come from an infested area (T Yaacobi, pers. comm., 2017)
- seeds are likely to be transported by mud on machinery. The fruits have hooked spines that can attach easily to clothing, boot or shoelaces. The plant forms near monospecific stands in habitats where people walk (e.g. natural areas and agricultural land, roadsides).

Impact

Effects on plants

Ambrosia confertiflora forms dense stands that can outcompete native plant species (Fig. 2) (J. M. Dufour-Dror, pers. comm. 2017). The plant has the ability to modify the vegetation dynamics (Israel Ministry of Environmental



Fig. 2 Infestation of the bank of the Alexander River in Israel. Photo courtesy of J. M. Dufour.

Protection, 2013). A monoculture can act to suppress understorey native plants, resulting in environmental impacts, in particular in humid habitats, in grasslands and in dry river beds (J. M. Dufour-Dror, pers. comm. 2017). In Israel, the plant spreads rapidly in wetlands and thrives along riverbanks, where it displaces native species and creates pure stands (J. M. Dufour-Dror, pers. comm. 2014). Species that have been shown to be displaced by *A. confertiflora* in Israel include *Bolboschoenus maritimus*, *Asparagus palaestinus*, *Polygonum salicifolium*, *Vicia galeata*, *Cyperus fuscus*, *Verbena officinalis*, *Ipomoea sagittata*, *Ranunculus sceleratus*, *Epilobium hirsutum*, *Pulicaria dysenterica* (T. Yaacobi, pers. comm. 2017).

In Australia, there are no known studies that have assessed the impact of the species on biological diversity, but the species is regarded as suppressing pasture plants such as *Chloris gayana* (Rhodes grass) (Australian Government, 2017).

Environmental and social impact

In Israel, *A. confertiflora* has been shown to have a significant impact on ecosystem services although a lot of the data for this is observational rather than from scientific studies. *Ambrosia confertiflora* is also a serious pest in cultivated fields, citrus groves and orchards in general and in vineyards (Israel Ministry of Environmental Protection, 2013). The plant competes for nutrients and can spread among cultivated fields and mixes with crops which cannot be harvested properly (Parsons & Cuthbertson, 2001). In particular, *A. confertiflora* has a major impact on organic farming as it cannot be effectively controlled by cultural and mechanical practices. It has caused notable damage to *Moringa* spp. fields in the Sharon region (J. M. Dufour-Dror, pers. comm. 2017). The burrs of the plant contaminate wool and can lower its value (Parsons & Cuthbertson, 2001).

Infestations in recreational sites may be problematic because of the unpleasant smell of the plant and mainly

because of possible dermatitis and allergies it causes. In addition, monocultures can prevent access to sites and rivers. The plant also invades gardens and parks and is reported to disrupt their maintenance (J. M. Dufour-Dror, pers. comm. 2017).

As with the other *Ambrosia* species, *A. confertiflora* produces a large amount of pollen which is considered to be severely allergenic (Yair, 2017), causing hay fever and contact dermatitis in susceptible people (Parsons & Cuthbertson, 2001).

Control

Ambrosia confertiflora is not effectively controlled by cultural practices. In fact, cultural practices can make the infestation worse by spreading pieces of the perennial root and stimulating development of root buds. Mechanical control can even exacerbate the intensity of the invasion, as the plant regenerates very quickly after mowing (J. M. Dufour-Dror, pers. comm. 2017). Uprooting may only be effective against very young plants, as otherwise the root remains in the soil and the plant regenerates.

In Israel, herbicides such as glyphosate, triclopyr and fluroxypyr are not very effective as the plant recovers quickly after the spraying (J. M. Dufour-Dror, pers. comm. 2017). In addition, these herbicides are not registered for use in wetlands and riparian habitats. A recent unpublished experiment in Israel suggests that imazapyr is effective in killing mature specimens of the burr ragweed (J. M. Dufour-Dror, pers. comm. 2017). However, herbicides have been recorded as providing satisfying results in controlling *A. confertiflora* in pastures in Australia.

An unpublished study conducted in Israel showed that restoration of natural vegetation along river banks with *Arundo donax* and *Arundo mediterranea* (Poaceae) successfully prevented the re-establishment of *A. confertiflora* in previously infested areas (J. M. Dufour-Dror, pers. comm. 2017).

Regulatory status

At present, within the EPPO region there are regulations specific to *A. confertiflora*. *Ambrosia confertiflora* was included on the EPPO List of Invasive Alien Plants in 2014. In 2016, the species was prioritized (along with 36 additional species from the EPPO List of Invasive Alien Plants and a recent horizon scanning study) for PRA within the LIFE-funded project ‘Mitigating the threat of invasive alien plants to the EU through PRA to support the Regulation 1143/2014’. *Ambrosia confertiflora* was one of 16 species identified as having a high priority for PRA (Branquart *et al.*, 2016; Tanner *et al.*, 2017). A subsequent PRA concluded that *A. confertiflora* had a high phytosanitary risk to the endangered area (EPPO, 2018) and was added to the EPPO A2 List of pests recommended for regulation. At the time of publishing, *A. confertiflora* is being

considered for inclusion on the list of (European) Union concern.

Ambrosia confertiflora is declared ‘Noxious’ in South Australia (proclaimed plant as *Ambrosia* spp.) (Natural Resources Management Act). In addition, in Australia, the species is a non-saleable weed.

Acknowledgements

This datasheet is an output of a DG Environment, LIFE funding, under the project LIFE15 PRE-FR 001: Mitigating the threat of invasive alien plants in the EU through pest risk analysis to support the EU Regulation 1143/2014. The datasheet was produced following an Expert Working Group (EWG) that risk analysed *A. confertiflora* for the EPPO region in September 2017. The composition of the EWG was as follows: M. Bouhache (Institut Agronomique et Vétérinaire, MA), D. Chapman (Centre for Ecology and Hydrology, GB), S. Follak (Austrian Agency for Health and Food Safety (AGES), AT), J. Lambrinos (Oregon State University, US), J. van Valkenburg (National Plant Protection Organization, NL), V. Visser (University of Cape Town, ZA), T. Yaacobi (Research and Consultancy in Applied Ecology, IL) and R. Tanner (EPPO).

References

- Australian Government (2017) Weeds in Australia, *Ambrosia confertiflora*. http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=15367 [accessed on 25 September 2017]
- Branquart E, Brundu G, Buholzer S, Chapman D, Ehret P, Fried G *et al.* (2016) A prioritization process for invasive alien plant species incorporating the requirements of the EU Regulation 1143/2014. *EPPO Bulletin* **47**, 603–617.
- Calflora (2017) *Ambrosia confertiflora*. https://www.calflora.org/cgi-bin/species_query.cgi?where-calrecnum=294 [accessed on 25 September 2017]
- Dufour-Dror JM (2012) *Alien invasive plants in Israel*, p. 213. The Middle East Nature Conservation Promotion Association, Ahva publishing, Jerusalem (IL).
- Dufour-Dror JM (2016) Improving the prevention of alien plant invasion in the EPPO region: the need to focus on highly invasive plants with (still) limited distribution—examples from Israel. *EPPO Bulletin* **46**(2), 341–347.
- Dufour-Dror JM & Fragman-Sapir O (2018) Alien plants species in natural and disturbed areas in Israel. In *Alien Invasive plants in Israel*, 2nd edn (ed. Dufour-Dror JM). Cana publishing, Jerusalem (IL), in press.
- Dufour-Dror JM & Yaacoby T (2013) *Ambrosia confertiflora*, *Burr ragweed – Factsheet*, p. 1–7. Israel Ministry of Environmental Protection, Israel Nature & Parks Authority, Israel Plant Protection & Inspection Services, Jerusalem.
- EPPO (2018) *Pest risk analysis for Ambrosia confertiflora EPPO*, Paris. <https://pra.eppo.int/> [accessed on 8 November 2018]
- Flora of North America (2017) *Ambrosia confertiflora*. http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=250066047 [accessed on 23 September 2017]

- Israel Ministry of Environmental Protection (2013) *Ambrosia confertiflora*, Burr Ragweed, Information booklet. p. 7. <http://www.sviva.gov.il/InfoServices/ReservoirInfo/DocLib2/Publications/P0701-P0800/P0707.pdf> [accessed on 23 September, 2017]
- Parsons WT & Cuthbertson EG (2001) *Noxious weeds of Australia*, 2nd edn, p. 698. CSIRO Publishing, Collingwood (AU).
- Richardson FJ, Richardson RG & Shepherd RCH (2011) *Weeds of the South-East – An identification guide for Australia*, 2nd edn, p. 546. R.G & F.J. Richardson, Meredith, Vic. (AU).
- Tanner R, Branquart E, Brundu G, Buholzer S, Chapman D, Ehret P *et al.* (2017) The prioritisation of a short list of alien plants for risk analysis within the framework of the Regulation (EU) No. 1143/2014. *NeoBiota* **35**, 87–118.
- Yaacoby T & Seplyarsky V (2011) *Epiblema strenuana* (Walker, 1863) (Lepidoptera: Tortricidae), a new species in Israel. *EPPO Bulletin* **41** (2), 243–246.
- Yair Y (2017) *Ambrosia* spp. In Israel, an invasive and allergenic weed. PhD thesis, Tel Aviv University.