

Data Sheets on Quarantine Pests

*Endocronartium harknessii***IDENTITY**

Name: *Endocronartium harknessii* (J.P. Moore) Y. Hiratsuka

Synonyms: *Cronartium harknessii* E. Meinecke
Peridermium harknessii J.P. Moore

Taxonomic position: Fungi: Basidiomycetes: Uredinales

Common names: Western gall rust, pine-pine gall rust (English)

Notes on taxonomy and nomenclature: In the past, a broad concept of *C. coleosporioides* (EPPO/CABI, 1996) was followed, by which it was considered to include a set of distinctive aecial stages: *Peridermium stalactiforme* (northern distribution), *P. filamentosum* (southern distribution) and *P. harknessii* (ubiquitous and autoecious) (Mordue & Gibson, 1978). However, Hiratsuka (1969) concluded that the aeciospores of *P. harknessii* function as teliospores and erected the new genus *Endocronartium* to accommodate such endocyclic life cycles. *E. harknessii* is thus now considered as a distinct species.

Bayer computer code: ENDCHA

EPPO A1 list: No. 11

EU Annex designation: I/A1 - as *Endocronartium* spp. (non-European)

HOSTS

The aecial hosts of *E. harknessii* in North America are two- and three-needled *Pinus* spp., of which the most important are jack pine (*P. banksiana*) across Canada, and lodgepole pine (*P. contorta*) and western yellow pine (*P. ponderosa*) in western Canada and USA. The European Scots pine (*P. sylvestris*), widely planted in North America, is susceptible. Other recorded aecial hosts include the locally less important far western species knobcone pine (*P. attenuata*), bishop pine (*P. muricata*) and Monterey pine (*P. radiata*). The European species Aleppo pine (*P. halepensis*), mountain pine (*P. mugo*) and Austrian pine (*P. nigra*) have been found to be infected in North America. In view of the fact that *P. contorta* is widely planted in northern and western Europe, *P. radiata* in western Europe and *P. ponderosa* to a certain extent in central Europe, and the previously mentioned European species are also susceptible, *E. harknessii* would certainly find hosts on which to establish in the EPPO region. Since it is, by nature, an autoecious short-cycle rust, there are no telial hosts (the aeciospores of *E. harknessii* function as teliospores, leading to reinfection of the original *Pinus* host). For more information, see Spaulding (1956, 1961), Boyce (1961), USDA (1963), Davidson & Prentice (1967), Peterson (1967), Hepting (1971), Ziller (1974).

GEOGRAPHICAL DISTRIBUTION

EPPO region: Absent.

North America: Canada (practically throughout - Alberta, British Columbia, Manitoba, New Brunswick, Nova Scotia, Northwest Territory, Ontario, Quebec, Saskatchewan, Yukon Territory), Mexico, USA (western and northern states Alaska, Arizona, California, Colorado, Idaho, Massachusetts, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New York, North Dakota, Oregon, Pennsylvania, Virginia, Washington).

Central America and Caribbean: Unconfirmed reports in Central America.

South America: Unconfirmed reports.

EU: Absent.

Distribution map: See CMI (1993, No. 449).

BIOLOGY

The life history of *E. harknessii* is remarkably simple: aecidioid teliospores (spores having the morphological characteristics of aeciospores but germinating and functioning like teliospores) develop in the spring on branch galls (rarely on stem cankers) of *Pinus* 2-4 years after infection; these spores are carried in air currents and reinfect *Pinus* directly. Once established, galls continue to produce spores each spring; invasion by secondary fungi usually results in death of the branch. The rust may continue to survive in cankers for up to 200 years, but produces few, if any, spores. Van der Kamp (1994) found that most infections of *P. contorta* by *E. harknessii* occurred within 2 m of the ground and disappeared as the branches concerned were progressively shed, without new infections appearing. For more information, see also Peterson (1960), Ziller (1967, 1974), Krebill (1970), Byler & Platt (1972), Peterson (1973).

DETECTION AND IDENTIFICATION

Symptoms

In spring, droplets of clear, viscous liquid are produced. These are followed by pale-yellow aecia, 1-8 mm in diameter, containing powdery yellow-orange spores. Each new infection is followed by the formation of a well delimited, spherical to oblong gall, up to 8 cm, sometimes accompanied by a small witches' broom. Small galls on twigs 1-2 years old are frequently pyriform. The bark sloughs off in large scales, finally exposing the smooth wood, with a collar of old bark standing out around the upper and lower ends of the gall. Galls form on branches and small stems of *Pinus* of all ages. For more information, see also Boyce (1961), USDA (1963), Hepting (1971), Ziller (1974), Sinclair *et al.* (1987).

Morphology

Teliospores subglobose to obovoid and ellipsoid, with colourless, coarsely verrucose walls which have a lateral smooth spot; 23-35 x 14-24 μm . They produce germ tubes (basidia) with one to four septa. See also Peterson (1960), Ziller (1967, 1974), Sinclair, *et al.* (1987). The fungus has been cultured (Allen *et al.*, 1988).

Detection and inspection methods

Isoenzyme and protein analysis of aeciospores can differentiate between *E. harknessii* and various *Cronartium* spp. (e.g. *C. quercuum*) in, for example, *P. banksiana* (Powers *et al.*, 1989; Tuskan & Walla, 1989).

MEANS OF MOVEMENT AND DISPERSAL

E. harknessii can be carried considerable distances as wind-borne aeciospores and can survive long periods in the airborne state (Chang & Blenis, 1989). More importantly, it can

also be carried to new areas on plants for planting of the coniferous aecial hosts, as has occurred in parts of the USA. Long incubation periods mean that latent infections easily go undetected unless post-entry quarantine is applied. There is no risk in the movement of *Pinus* seeds or pollen.

PEST SIGNIFICANCE

Economic impact

E. harknessii is damaging in its effect on form, lumber content and growth rates of *Pinus*, and kills individual trees, although it is not known to wipe out whole stands. Galls on the main stem of young trees can lead to the death of the tree, but galls on branches of older trees cause little loss (Gross, 1983). Severe outbreaks on seedlings of *P. ponderosa* and *P. contorta* have been recorded in north-west Canada as well as serious damage on young natural stands of *P. banksiana* and *P. sylvestris* plantations in Quebec. Damage in commercial stands occurs chiefly to *P. contorta* in the Rocky Mountains (Ziller, 1967, 1974). *P. sylvestris* has recently been reported to be much less susceptible than *P. contorta* (Van der Kamp, 1989). When compared with "blister rusts" (e.g. *C. comandrae*) on *P. banksiana* and *P. contorta* in northern Canada, *E. harknessii* was found not to be so aggressive on dominant trees, but to attack intermediate or suppressed trees which would disappear in any case (Hiratsuka *et al.*, 1988).

Control

The autoecious nature of the rust makes control difficult. Removal of infected trees may be economical. Nurseries should be well away from infection foci. Several non-rust fungi have been reported in association with and acting as natural biological control agents of *E. harknessii*. *Scytalidium uredinicola* has attracted particular interest, and produces a spore germination inhibitor of *E. harknessii* (Cunningham & Pickard, 1985). Resistance to *E. harknessii* also exists (Yanchuk *et al.*, 1988; Burnes *et al.*, 1989).

Phytosanitary risk

E. harknessii is an EPPO A1 quarantine pest (OEPP/EPPO, 1979), and is also a quarantine pest for IAPSC. It is considered a major quarantine pest for Australia (Commonwealth Department of Health, 1982). The potential danger of *E. harknessii* to commercially important *Pinus* in the EPPO region, especially the native species *P. sylvestris*, but also other native species and introduced North American species, can clearly be seen, in view of its considerable importance in North America, the difficulties of control and the fact that the fungus can spread from tree to tree, without an alternate host.

PHYTOSANITARY MEASURES

Since symptoms may not be apparent for several years after infection, the only practical safeguard is to prohibit entry of the *Pinus* species which are hosts (see Hosts) from countries where *E. harknessii* occurs (OEPP/EPPO, 1990). Bark and wood of host *Pinus* should have been appropriately treated (heat-treated, fermented, kiln-dried; EPPO quarantine procedures are in preparation).

BIBLIOGRAPHY

- Allen, E.A.; Blenis, P.V.; Hiratsuka, Y. (1988) Axenic culture of *Endocronartium harknessii*. *Mycologia* **80**, 120-123.
- Boyce, J.S. (1961) *Forest pathology* (3rd edition), pp. 201-217. McGraw-Hill Book Co., New York, USA.

- Burnes, T.A.; Blanchette, R.A.; Stewart, W.K.; Mohn, C.A. (1989) Screening jack pine seedlings for resistance to *Cronartium quercuum* f.sp. *banksianae* and *Endocronartium harknessii*. *Canadian Journal of Forest Research* **19**, 1642-1644.
- Byler, J.W.; Platt, W.D. (1972) Cone infection by *Peridermium harknessii*. *Canadian Journal of Botany* **50**, 1429-1430.
- Chang, K.F.; Blenis, P.V. (1989) Survival of *Endocronartium harknessii* teliospores in a simulated airborne state. *Canadian Journal of Botany* **67**, 928-932.
- CMI (1993) *Distribution Maps of Plant Diseases* No. 449 (edition 3). CAB International, Wallingford, UK.
- Commonwealth Department of Health (1982) Western gall rust. *Plant Quarantine Leaflet* No. 28. Commonwealth Department of Health, Canberra, Australia.
- Cunningham, J.E.; Pickard, M.A. (1985) Maltol, a metabolite of *Scytalidium uredinicola* which inhibits spore germination of *Endocronartium harknessii*, the western gall rust. *Canadian Journal of Microbiology* **31**, 1051-1055.
- Davidson, A.G.; Prentice, R.M. (1967) Important forest insects and diseases of mutual concern to Canada, the United States and Mexico. *Department of Forest and Rural Development, Canada Publication* No. 1180.
- EPPO/CABI (1996) *Cronartium coleosporioides*. In: *Quarantine pests for Europe*. 2nd edition (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Holderness, M.). CAB INTERNATIONAL, Wallingford, UK.
- Gross, H.L. (1983) Negligible cull and growth loss of jack pine associated with globose gall rust. *Forestry Chronicle* **59**, 308-311.
- Hepting, G.H. (1971) *Diseases of forest and shade trees of the United States*. Agricultural Handbook, Forest Service, US Department of Agriculture No. 386, pp. 287-370.
- Hiratsuka, Y. (1969) *Endocronartium*, a new genus for autoecious pine stem rusts. *Canadian Journal of Botany* **47**, 1493-1495.
- Hiratsuka, Y.; Powell, J.M.; Van Sickle, G.A. (1988) Impact of pine stem rusts of hard pines in Alberta and the Northwest Territories. *Information-Report - Northern Forestry Centre, Canadian Forestry Service* No. NOR-X-299.
- Krebill, R.G. (1970) Autoecious gall rusts of pines in South Michigan and New York. *Plant Disease Reporter* **54**, 853-855.
- Mordue, J.E.M.; Gibson, I.A.S. (1978) *Cronartium coleosporioides*. *CMI Descriptions of Pathogenic Fungi and Bacteria* No. 577. CAB International, Wallingford, UK.
- OEPP/EPPO (1979) Data sheets on quarantine organisms No. 11, *Endocronartium harknessii*. *Bulletin OEPP/EPPO* **9** (2).
- OEPP/EPPO (1990) Specific quarantine requirements. *EPPO Technical Documents* No. 1008.
- Peterson, G.W. (1973) Dispersal of aeciospores of *Peridermium harknessii* in central Nebraska. *Phytopathology* **63**, 170-172.
- Peterson, R.S. (1960) Western gall rust on hard pines. *Forest Pest Leaflet, Forest Service, US Department of Agriculture* No. 50, 8 pp.
- Peterson, R.S. (1967) The *Peridermium* species on pine stems. *Bulletin of the Torrey Botanical Club* **94**, 511-542.
- Powers, H.R.; Lin, D.; Hubbes, M. (1989) Interspecific and intraspecific differentiation within the genus *Cronartium* by isozyme and protein pattern analysis. *Plant Disease* **73**, 691-694.
- Sinclair, W.A.; Lyon, H.H.; Johnson, W.T. (1987) In: *Diseases of trees and shrubs*, 574 pp. Comstock Publishing Associates, Ithaca, USA.
- Spaulding, P. (1956) Diseases of North American forest trees planted abroad. An annotated list. *Agricultural Handbook, Forest Service, US Department of Agriculture* No. 100, p. 11.
- Spaulding, P. (1961) Foreign diseases of forest trees of the world. An annotated list. *Agricultural Handbook, Forest Service, US Department of Agriculture* No. 197, pp. 74, 183.
- Tuskan, G.A.; Walla, J.A. (1989) Isozyme characterization of *Peridermium harknessii* and *Cronartium quercuum* f.sp. *banksianae* with starch gel electrophoresis. *Phytopathology* **79**, 444-448.
- USDA (1963) Internationally dangerous forest tree diseases. *Miscellaneous Publications, Forest Service, US Department of Agriculture* No. 939, pp. 54, 56-57, 73-74, 92-96.

- Van der Kamp, B.J. (1989) The relative susceptibility of Scots and lodgepole pine to western gall rust. *European Journal of Forest Pathology* **19**, 274-280.
- Van der Kamp, B.J. (1994) Lodgepole pine stem diseases and management of stand density in the British Columbia interior. *Forestry Chronicle* **70**, 773-779.
- Yanchuk, A.D.; Yeh, F.C.; Dancik, B.P. (1988) Variation of stem rust resistance in a lodgepole pine provenance-family plantation. *Forest Science* **34**, 1067-1075.
- Ziller, W.G. (1967) Western gall rust. In: *Important forest insects and diseases of mutual concern to Canada, the United States and Mexico* (Ed. by Davidson, A.G.; Prentice, R.M.). *Department of Forest and Rural Development, Canada Publication No. 1180*, pp. 53-54.
- Ziller, W.G. (1974) The tree rusts of Western Canada. *Forest Service, British Columbia, Canada Publication No. 1329*, pp. 78-100.