

# EPPO Datasheet: *Diaphorina citri*

Last updated: 2020-09-03

## IDENTITY

**Preferred name:** *Diaphorina citri*

**Authority:** Kuwayana

**Taxonomic position:** Animalia: Arthropoda: Hexapoda: Insecta: Hemiptera: Sternorrhyncha: Liviidae

**Other scientific names:** *Euphalerus citri* (Kuwayana)

**Common names:** Asian citrus psyllid, citrus psylla, citrus psyllid  
[view more common names online...](#)

**EPPO Categorization:** A1 list

[view more categorizations online...](#)

**EU Categorization:** A1 Quarantine pest (Annex II A)

**EPPO Code:** DIAACI



[more photos...](#)

## Notes on taxonomy and nomenclature

*Diaphorina citri* was recently moved from the family Psyllidae to the family Liviidae. Literature prior to 2015 will probably allocate this species in Psyllidae.


## HOSTS

*D. citri* is confined to Rutaceae, occurring on wild hosts and on cultivated *Citrus*, especially grapefruit (*Citrus paradisi*), lemons (*C. limon*) and limes (*C. aurantiifolia*). *Murraya paniculata*, a rutaceous plant often used for hedges, is a preferred host. Within the EPPO region, host species are generally confined to countries surrounding the Mediterranean Sea.

**Host list:** *Aegle marmelos*, *Afraegle paniculata*, *Archidendron lucidum*, *Atalantia buxifolia*, *Atalantia*, *Balsamocitrus dawei*, *Casimiroa edulis*, *Citroncirus webberi*, *Citroncirus*, *Citrus amblycarpa*, *Citrus aurantiifolia*, *Citrus aurantium*, *Citrus australasica*, *Citrus australis*, *Citrus halimii*, *Citrus hassaku*, *Citrus hystrix*, *Citrus inodora*, *Citrus jambhiri*, *Citrus latipes*, *Citrus limettioides*, *Citrus limon*, *Citrus macrophylla*, *Citrus maxima*, *Citrus medica*, *Citrus paradisi*, *Citrus reshni*, *Citrus reticulata*, *Citrus sinensis*, *Citrus sunki*, *Citrus taiwanica*, *Citrus volkameriana*, *Citrus webberi*, *Citrus x limonia*, *Citrus x nobilis*, *Citrus*, *Clausena anisum-olens*, *Clausena excavata*, *Clausena harmandiana*, *Clausena indica*, *Clausena lansium*, *Cordia myxa*, *Eremocitrus glauca*, *Ficus carica*, *Fortunella japonica*, *Fortunella sp.*, *Fortunella*, *Glycosmis pentaphylla*, *Limonia acidissima*, *Merrillia caloxylon*, *Murraya koenigii*, *Murraya paniculata*, *Poncirus trifoliata*, *Rutaceae*, *Swinglea glutinosa*, *Toddalia asiatica*, *Triphasia trifolia*, *Vepris lanceolata*, *Zanthoxylum ailanthoides*, *x Citrofortunella microcarpa*, *x Citrofortunella sp.*

## GEOGRAPHICAL DISTRIBUTION

The distribution of *D. citri* is wider than that of the causal agent of citrus huanglongbing (HLB) disease originally associated with this vector, '[Candidatus Liberibacter asiaticus](#)'. *D. citri* occurs in Afghanistan, some states of Brazil, Macau, Myanmar, and Singapore, where the associated bacterium has not yet been recorded.



**Africa:** Ethiopia, Kenya, Mauritius, Nigeria, Reunion, Tanzania

**Asia:** Afghanistan, Bangladesh, Bhutan, Cambodia, China (Aomen (Macau), Fujian, Guangdong, Guangxi, Guizhou, Hainan, Henan, Hunan, Jiangxi, Sichuan, Xianggang (Hong Kong), Yunnan, Zhejiang), East Timor, India (Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Delhi, Gujarat, Haryana, Himachal Pradesh, Jammu & Kashmir, Karnataka, Kerala, Lakshadweep, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Punjab, Rajasthan, Sikkim, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal), Indonesia (Java, Maluku, Nusa Tenggara, Sumatra), Iran, Japan (Kyushu, Ryukyu Archipelago), Lao, Malaysia (Sabah, West), Maldives, Myanmar, Nepal, Oman, Pakistan, Philippines, Saudi Arabia, Singapore, Sri Lanka, Taiwan, Thailand, United Arab Emirates, Viet Nam, Yemen

**North America:** Mexico, United States of America (Alabama, Arizona, California, Florida, Georgia, Hawaii, Louisiana, Mississippi, South Carolina, Texas)

**Central America and Caribbean:** Antigua and Barbuda, Bahamas, Barbados, Belize, Cayman Islands, Costa Rica, Cuba, Dominica, Dominican Republic, Guadeloupe, Haiti, Jamaica, Martinique, Puerto Rico, Saint Lucia, St Vincent and the Grenadines, Virgin Islands (US)

**South America:** Argentina, Brazil (Amazonas, Bahia, Ceara, Para, Pernambuco, Rio de Janeiro, Santa Catarina, Sao Paulo), Colombia, Paraguay, Uruguay, Venezuela

**Oceania:** American Samoa, Guam, Northern Mariana Islands, Papua New Guinea

## BIOLOGY

*D. citri* has a short life cycle and high fecundity. Its optimal developmental temperature ranges from 25-28°C (Liu and Tsai, 2000), and it is, therefore, best adapted to tropical and subtropical conditions, although mean temperatures above 30°C reduce its survival and fertility. Pairing starts soon after emergence, the insects being most active in spring and summer, once mean temperatures are above 12°C (Udell *et al.*, 2017). Eggs are laid inside half-folded leaves in buds, in leaf axils or other suitable places on the young tender parts of the tree (shoots or flushes). Females have a pre-oviposition period of about 10 days at their optimal temperature, and can lay up to 800 eggs during their lifetime (Liu and Tsai, 2000). Eggs hatch within 3-10 days (3 days at 28°C and 10 days at 15°C) and nymphs pass through five instars in 11-40 days. The complete life cycle takes 14-50 days, with up to 9-14 overlapping generations per year at mean temperatures of 20-25°C. However, as the eggs are laid exclusively on young flushes and nymphs can only develop on tender plant tissue, the number of generations per year is limited by the sprouting activity of citrus trees, and population fluctuations are closely correlated with tree phenology during the growing season (Udell *et al.*, 2017). *D. citri* overwinters as an adult, which can live for up to six months. Adults are highly active and jump at the slightest disturbance. Nymphs will move away when disturbed, but normally lead a sedentary existence, clustered in groups.

*D. citri* is known to be the most efficient vector of '*Ca. Liberibacter asiaticus*', the most aggressive causal agent of Huanglongbing disease. Together, these two organisms constitute the most destructive citrus pathosystem worldwide

(Gottwald, 2010). *D. citri* can also transmit ‘*Candidatus Liberibacter africanus*’ and ‘*Candidatus Liberibacter americanus*’. In areas in which these three bacteria species coexist, *D. citri* can transmit them indiscriminately (Ajene *et al.*, 2020; Gottwald, 2010).

## **DETECTION AND IDENTIFICATION**

### **Symptoms**

High densities of *D. citri* nymphs feeding on tender shoots can stunt and twist them. Lateral leaf notching is a characteristic form of damage associated with this insect (Aubert, 1987). However, at low insect densities, this symptom may go unnoticed. Both adults and immature insects secrete a semi-solid honeydew. Sooty mold may occur in the presence of high psyllid densities and humid environmental conditions.

### **Morphology**

#### **Eggs**

Yellow, almond-shaped and tapering at the distal end; 0.01-0.15 mm.

#### **Nymph**

Light-yellow to dark-brown or green, bearing well-developed wing pads.

#### **Adult**

Adults are about 2-4 mm long, with a yellowish-brown body, greenish-brown or pinkish-brown abdomen, and greyish-brown legs. Males are smaller than females. The wings are transparent, mottled with white and brown spots and a broad, beige, band is present at the periphery of the distal part, slightly interrupted near the apex. The terminal segments of the antennae are black. Two darker segments are also found in the middle of the antennae.

The EPPO Diagnostic Protocol on *D. citri* provides guidance on how to detect and identify the pest (EPPO Standard PM 7/52).

### **Detection and inspection methods**

Several sampling methods have been proposed and tested for the detection and monitoring of *D. citri*. The sampling methods recommended depend on the goals of the monitoring program. For early detection, suction sampling devices for the capture of adults, and yellow sticky traps are mostly recommended. For regular *D. citri* management actions, the stem tap sampling of adults provides reliable information rapidly. The visual sampling of nymphs in tender shoots during the major citrus sprouting periods of the tree growing season is recommended for determinations of the number of *D. citri* generations (Monzo and Stansly, 2020). Detailed protocols for surveillance, sampling and detection are indicated in the EPPO Standard PM 9/27 (2020) and in the EFSA pest survey card (EFSA, 2019).

## **PATHWAYS FOR MOVEMENT**

*D. citri* has a substantial flight capacity. It can disperse locally over distances of at least 2 km, within 12 days, when food and oviposition resources (tender citrus shoots) are scarce (Lewis-Rosenblum *et al.*, 2015). Eggs and nymphs can be carried over longer distances on citrus material (budwood, grafted trees, rootstock seedlings) from infected areas. Trailers transporting oranges from groves to packing houses have also been recognized as a source of adult vector dispersal (Halbert *et al.* 2010). Adults and, 5th or 6th-instar nymphs can transmit '*Ca. Liberibacter asiaticus*' to citrus plants. However, cleaned fruit, that has been washed and is without leaves at the end of the packing process is not considered to pose a risk. The rutaceous plant *Murraya paniculata*, frequently used as an ornamental bush or hedge, is one of the preferred hosts of *D. citri*. This plant can carry *D. citri* eggs or nymphs, and its introduction into disease- and vector-free regions creates a risk of introduction of huanglongbing or its vector.

## PEST SIGNIFICANCE

### Economic impact

The economic impact of *D. citri* results principally from its role as a vector of huanglongbing, the most damaging citrus disease in the world (Gottwald, 2010). The presence of this pathosystem systematically drastically affects the citrus industry (Hodges and Spreens, 2012), and significantly impairs integrated pest management strategies (Grafton-Cardwell *et al.*, 2013). This pathosystem greatly decreases tree productivity, increases management costs considerably, and may also have deleterious effects on fruit quality (Tansey *et al.*, 2017). Intensive management programs for this pathosystem also have a negative impact on biological control processes for this crop, with important economic consequences (Monzo and Stansly, 2020a). In addition, direct feeding by *D. citri* causes leaf curling and notching. Under heavy infestations, sprouting shoots die, resulting in blossom and fruitlet drop.

### Control

There are no curative treatments for huanglongbing, and there are no tolerant or resistant plants. The management of this pathosystem is therefore entirely dependent on efficient vector control (Grafton-Cardwell *et al.*, 2013). Insecticide-based strategies are the most effective, but must be compatible with biological control (Monzo *et al.*, 2014; Qureshi *et al.*, 2009). Broad-spectrum insecticide use in the winter is recommended, as only adult psyllids are found on citrus plants at this time of the year (Qureshi and Stansly, 2010). More selective active ingredients must be used during the tree growing season, when biological control is more relevant (Monzo *et al.*, 2014; Qureshi and Stansly, 2009). Due to the high frequency of insecticide applications against this vector in huanglongbing management programs, the rotation of modes of action is essential, to reduce the risk of resistance (Tiwari *et al.*, 2011). Economic thresholds have also been proposed, for spraying based on *D. citri* abundance, as a means of reducing the number of applications (Monzo and Stansly 2017). Biological control alone is not sufficient to prevent the spread of huanglongbing, but it can help to reduce the frequency of applications in commercial groves and is the most efficient management strategy for citrus trees in non-commercial citrus growing areas, such as those in urban gardens (Kistner *et al.* 2016; Monzo and Stansly, 2020a). The parasitoid *Tamarixia radiata* has been introduced into several citrus-growing regions worldwide, with various degrees of success (Chen and Stansly, 2014). Conservation biological control of naturally occurring predators can also greatly reduce the size of *D. citri* populations during the citrus growing season (Qureshi and Stansly, 2009; Monzo *et al.* 2014).

### Phytosanitary risk

*D. citri* could probably establish itself and spread in Mediterranean countries without difficulty. The presence of *D. citri* would greatly increase the risk of introduction and spread of huanglongbing. However, in addition to its role in spreading huanglongbing, this psyllid has a significant potential for damage in its own right. Biological control may be possible, but there is no guarantee that it could keep populations at sufficiently low levels to prevent transmission of huanglongbing.

## PHYTOSANITARY MEASURES

Considering the severity of huanglongbing, EPPO has recommended to prohibit the importation of citrus plants for planting and cut branches or buds of citrus from areas or countries where citrus huanglongbing (or either of its

vectors) are present. In the EU territory, it is also forbidden to import fruit from third countries with their peduncles and leaves. In disease free countries as those of the Mediterranean area, awareness, monitoring, surveillance, pest risk assessment, quarantine measures and action plans are advised (Duran-Vila *et al.*, 2014; Siverio *et al.*, 2017). Procedures for official control with the aim of detecting, containing and eradicating huanglongbing and its vectors are provided in the EPPO Standard PM 9/27 (EPPO, 2020). As surveys should be carried out in all the EU member countries, a pest survey card was prepared by the European Food Safety Authority (EFSA, 2019) to assist EU Member States in planning their huanglongbing annual survey activities.

## REFERENCES

- Ajene IJ, Khami FM, van Asch B, Pietersen G, Seid N, Rwomushana I, Ombura FLO, Momanyi G, Finyange P, Rasowo BA, Tanga CM, Mohammed S, Ekesi S (2020) Distribution of *Candidatus Liberibacter* species in Eastern Africa, and the first report of *Candidatus Liberibacter asiaticus* in Kenya. *Scientific Reports* **10**, 3919 <https://doi.org/10.1038/s41598-020-60712-0>
- Aubert B (1987) *Trioza erytrae* Del Guercio and *Diaphorina citri* Kuwayama (Homoptera: Psylloidea), the two vectors of citrus greening disease: Biological aspects and possible control strategies. *Fruits*, **42**, 149-162.
- Chen X, Stansly PA (2014). Biology of *Tamarixia radiata* (Hymenoptera: Eulophidae), parasitoid of the citrus greening disease vector *Diaphorina citri* (Hemiptera: Psylloidea): a mini review. *Florida Entomologist*, **97**, 1404-1413.
- Duran-Vila, N, Janse, JD, Foissac, X, Melgarejo, P, Bove, JM (2014). Addressing the threat of Huanglongbing in the Mediterranean region: a challenge to save the citrus industry. *Journal of Plant Pathology*, **96**, 4-3.
- EFSA (2019) Parnell S, Camilleri M, Diakaki M, Schrader G & Vos S. Pest survey card on Huanglongbing and its vectors. *EFSA Supporting publication* 2019, EN-1574. <https://doi.org/doi:10.2903/sp.efsa.2019.EN-1574>
- EPPO (2020) PM 9/27 (1) '*Candidatus Liberibacter*' species that are casual agents of Huanglongbing disease of citrus and their vectors: procedures for official control. *EPPO Bulletin*, **50**, 122-141.
- Gottwald TR (2010). Current epidemiological understanding of citrus huanglongbing. *Annual Review of Phytopathology*, **48**, 119-139.
- Grafton-Cardwell EE, Stelinski LL, Stansly PA (2013) Biology and management of Asian citrus psyllid, vector of the huanglongbing pathogens. *Annual Review of Entomology*, **58**, 413-432.
- Halbert SE, Manjunath KL, Ramadugu C, Brodie MW, Webb SE, Lee RF (2010) Trailers transporting oranges to processing plants move Asian citrus psyllids. *Florida Entomologist*, **93**, 33-38.
- Hodges AW, Spreen TH. Economic impacts of citrus greening (HLB) in Florida, 2006/07-2010/11. *EDIS*. 2012; **FE903**:1-6.
- Kistner EJ, Amrich R, Castillo M, Strode V, Hoddle MS (2016) Phenology of Asian citrus psyllid (Hemiptera: Liviidae), with special reference to biological control by *Tamarixia radiata*, in the residential landscape of southern California. *Journal of Economic Entomology*, **109**, 1047-1057.
- Lewis-Rosenblum H, Martini X, Tiwari S, Stelinski LL (2015) Seasonal movement patterns and long-range dispersal of Asian citrus psyllid in Florida citrus. *Journal of Economic Entomology*, **108**, 3-10.
- Liu YH, Tsai JH (2000) Effects of temperature on biology and life table parameters of the Asian citrus psyllid, *Diaphorina citri* Kuwayama (Homoptera: Psyllidae). *Annals of Applied Biology*, **137**, 201-206.
- Monzo C, Qureshi JA, Stansly PA (2014) Insecticide sprays, natural enemy assemblages and predation on Asian citrus psyllid, *Diaphorina citri* (Hemiptera: Psyllidae). *Bulletin of Entomological Research*, **104**, 576-585.
- Monzo C, Stansly PA (2017) Economic injury levels for Asian citrus psyllid control in process oranges from mature

trees with high incidence of huanglongbing. *PLoS One*, **12**. <https://doi.org/10.1371/journal.pone.0175333>

Monzó C, Stansly PA (2020a) Economic value of conservation biological control for management of the Asian citrus psyllid, vector of citrus Huanglongbing disease. *Pest Management Science*, **76**, 1691-1698.

Monzo C, Stansly PA (2020b) Sampling and economic thresholds for Asian citrus psyllid. In Asian Citrus Psyllid. Biology, Ecology and Management of the Huanglongbing Vector. JA Qureshi and PA Stansly editors. CABI, 352 pp.

Qureshi JA, Stansly PA (2009) Exclusion techniques reveal significant biotic mortality suffered by Asian citrus psyllid *Diaphorina citri* (Hemiptera: Psyllidae) populations in Florida citrus. *Biological Control*, **50**, 129-136.

Qureshi JA, Stansly PA (2010) Dormant season foliar sprays of broad-spectrum insecticides: An effective component of integrated management for *Diaphorina citri* (Hemiptera: Psyllidae) in citrus orchards. *Crop Protection*, **29**, 860-866.

Qureshi JA, Rogers ME, Hall DG, Stansly PA (2009) Incidence of invasive *Diaphorina citri* (Hemiptera: Psyllidae) and its introduced parasitoid *Tamarixia radiata* (Hymenoptera: Eulophidae) in Florida citrus. *Journal of Economic Entomology*, **102**, 247-256.

Siverio F, Marcos-Noales E, Bertolini E, Teresani G, Penalver J, Mansilla P, Aguin O, Perez-Otero R, Abelleira A, Guerrero-Garcia J, Hernandez E, Cambra M, Milagros-Lopez M (2017) Survey of huanglongbing associated to ‘*Candidatus Liberibacter*’ species in Spain: analyses of citrus plants and *Trioza erytreae*. *Phytopathologia Mediterranea* **56**, 98-110.

Tansey JA, Vanaclocha P, Monzo C, Jones M, Stansly PA (2017) Costs and benefits of insecticide and foliar nutrient applications to huanglongbing?infected citrus trees. *Pest Management Science*, **73**, 904-916.

Tiwari S, Mann RS, Rogers ME, Stelinski LL (2011). Insecticide resistance in field populations of Asian citrus psyllid in Florida. *Pest Management Science*, **67**, 1258-1268.

Udell BJ, Monzo C, Paris TM, Allan SA, Stansly PA (2017) Influence of limiting and regulating factors on populations of Asian citrus psyllid and the risk of insect and disease outbreaks. *Annals of Applied Biology*, **171**, 70-88.

## ACKNOWLEDGEMENTS

This datasheet was extensively revised in 2020 by Cesar Monzo from Instituto Valenciano de Investigaciones Agrarias (IVIA), Moncada, Valencia, Spain. His valuable contribution is gratefully acknowledged.

## How to cite this datasheet?

EPPO (2020) *Diaphorina citri*. EPPO datasheets on pests recommended for regulation. Available online. <https://gd.eppo.int>

## Datasheet history

This datasheet was first published in the EPPO Bulletin in 1988 and revised in the two editions of 'Quarantine Pests for Europe' in 1992 and 1997 as well as in 2020. It is now maintained in an electronic format in the EPPO Global Database. The sections on 'Identity', 'Hosts', and 'Geographical distribution' are automatically updated from the database. For other sections, the date of last revision is indicated on the right.

CABI/EPPO (1992/1997) *Quarantine Pests for Europe (1<sup>st</sup> and 2<sup>nd</sup> edition)*. CABI, Wallingford (GB).

EPPO (1988) Data sheets on quarantine organisms No. 151, Citrus greening bacterium and its vectors *Diaphorina citri* & *Trioza erytreae*. *Bulletin OEPP/EPPO Bulletin* 18, 497-507.